





Published:


- with international search report
- before the expiration of the time limit for amending the claims and to be republished in the event of receipt of amendments

For two-letter codes and other abbreviations, refer to the "Guidance Notes on Codes and Abbreviations" appearing at the beginning of each regular issue of the PCT Gazette.

HIV REPLICATION INHIBITING PYRIMIDINES AND TRIAZINES

- The present invention is concerned with pyrimidine derivatives having HIV (Human Immunodeficiency Virus) replication inhibiting properties. The invention further relates to methods for their preparation and pharmaceutical compositions comprising them. The invention also relates to the use of said derivatives for the manufacture of a medicament for the prevention or the treatment of HIV infection.
- The present invention is aimed at providing particular novel series of pyrimidine derivatives having HIV replication properties. WO 99/50250, WO 00/27825 and WO 01/85700 disclose certain substituted aminopyrimidines and WO 99/50256 and EP-834 507 disclose aminotriazines having HIV replication inhibiting properties.
- The compounds of the invention differ from the prior art compounds in structure, pharmacological activity and/or pharmacological potency. It has been found that the compounds of the invention not only act favorably in terms of their capability to inhibit the replication of Human Immunodeficiency Virus (HIV), but also by their improved ability to inhibit the replication of mutant strains, in particular strains which have become resistant to commercially available drugs (so-called drug or multi-drug resistant HIV strains).

Thus in one aspect, the present invention concerns a compound of formula



- a *N*-oxide, a pharmaceutically acceptable addition salt, a quaternary amine or a stereochemically isomeric form thereof, wherein
- $-a^1=a^2-a^3=a^4-$ represents a bivalent radical of formula
- (a-1); $-\text{CH}=\text{CH}-\text{CH}=\text{CH}-$
- (a-2); $-\text{N}=\text{CH}-\text{CH}=\text{CH}-$
- (a-3); $-\text{N}=\text{CH}-\text{N}=\text{CH}-$
- (a-4); $-\text{N}=\text{CH}-\text{CH}=\text{N}-$
- (a-5); $-\text{N}=\text{N}-\text{CH}=\text{CH}-$
- $-b^1-b^2-b^3-$ represents a bivalent radical of formula
- (b-1); $-\text{CH}_2-\text{CH}_2-\text{CH}_2-$
- n is 0, 1, 2, 3 or 4; and in case $-a^1=a^2-a^3=a^4-$ is (a-1), then n may also be 5;

-2-


m is 0, 1, 2, 3;

q is 0, 1 or 2;

p is 1 or 2;

R^1 is hydrogen; aryl; formyl; C_{1-6} alkylcarbonyl; C_{1-6} alkyl; C_{1-6} alkyloxycarbonyl; C_{1-6} alkyl substituted with formyl, C_{1-6} alkylcarbonyl, C_{1-6} alkyloxycarbonyl, C_{1-6} alkylcarbonyloxy; C_{1-6} alkyloxy C_{1-6} alkylcarbonyl substituted with C_{1-6} alkyloxycarbonyl;

each R^2 independently is hydroxy, halo, C_{1-6} alkyl optionally substituted with cyano or with $-C(=O)R^6$, C_{3-7} cycloalkyl, C_{2-6} alkenyl optionally substituted with one or more halogen atoms or cyano, C_{2-6} alkynyl optionally substituted with one or more halogen atoms or cyano, C_{1-6} alkyloxycarbonyl, carboxyl, cyano, nitro, $NR^{13}R^{14}$, polyhalomethyl, polyhalomethylthio, $-S(=O)_pR^6$, $-NH-S(=O)_pR^6$, $-C(=O)R^6$, $-NHC(=O)H$, $-C(=O)NHNH_2$, $-NHC(=O)R^6$, $-C(=NH)R^6$ or a radical of formula



wherein each A_i independently is N, CH or CR^6 ; and

A_2 is NH, O, S or NR^6 ;

X_1 is $-NR^5$ -, $-NH-NH$ -, $-N=N$ -, $-O$ -, $-C(=O)$ -, C_{1-4} alkanediyl, $-CHOH$ -, $-S$ -, $-S(=O)_p$ -, $-NR^{13}-C(=O)$ -, $-C(=O)-NR^{13}$ -, $-X_2-C_{1-4}$ alkanediyl- or $-C_{1-4}$ alkanediyl- X_2 -;

X_2 is $-NR^5$ -, $-NH-NH$ -, $-N=N$ -, $-O$ -, $-C(=O)$ -, $-CHOH$ -, $-S$ -, $-S(=O)_p$ -,

R^3 is hydrogen, halo, C_{1-6} alkyl, $NR^{13}R^{14}$, $-C(=O)-NR^{13}R^{14}$, $-C(=O)-R^{15}$, $-CH=N-NH-C(=O)-R^{16}$, $-C(=N-O-R^8)-C_{1-4}$ alkyl, R^7 or $-X_3-R^7$; or C_{1-6} alkyl substituted with one or more substituents each independently selected from halo, hydroxy, cyano, NR^9R^{10} , $-C(=O)-NR^9R^{10}$, $-C(=O)-C_{1-6}$ alkyl or R^7 , and in addition to said list of substituents, two geminal hydrogen atoms of said C_{1-6} alkyl may also be replaced by a C_{2-3} alkanediyl thus forming a spiro ring; C_{1-6} alkyloxy C_{1-6} alkyl optionally substituted with one or more substituents each independently selected from hydroxy, cyano, NR^9R^{10} , $-C(=O)-NR^9R^{10}$, $-C(=O)-C_{1-6}$ alkyl or R^7 ; C_{2-6} alkenyl substituted with one or more substituents each independently selected from halo, hydroxy, cyano, NR^9R^{10} , $-C(=O)-NR^9R^{10}$, $-C(=O)-C_{1-6}$ alkyl or R^7 ; C_{2-6} alkynyl substituted with one or more substituents each independently selected from halo, hydroxy, cyano, NR^9R^{10} , $-C(=O)-NR^9R^{10}$, $-C(=O)-C_{1-6}$ alkyl or R^7 ;





X_3 is $-NR^5$ -, $-NH-NH$ -, $-N=N$ -, $-O$ -, $-C(=O)$ -, $-S$ -, $-S(=O)_p$ -, $-X_2-C_{1-4}$ alkanediyl-, $-C_{1-4}$ alkanediyl- X_{2a} -, $-C_{1-4}$ alkanediyl- $X_{2b}-C_{1-4}$ alkanediyl-, $-C(=N-OR^8)-C_{1-4}$ alkanediyl-;

with X_{2a} being $-NH-NH$ -, $-N=N$ -, $-O$ -, $-C(=O)$ -, $-S$ -, $-S(=O)_p$ -; and
with X_{2b} being $-NH-NH$ -, $-N=N$ -, $-C(=O)$ -, $-S$ -, $-S(=O)_p$ -;

-3-

- R^4 is halo, hydroxy, C_{1-6} alkyl, C_{2-6} alkenyl, C_{2-6} alkynyl, C_{3-7} cycloalkyl, C_{1-6} alkyloxy, cyano, nitro, polyhalo C_{1-6} alkyl, polyhalo C_{1-6} alkyloxy, $-C(=O)-NR^{13}R^{14}$, C_{1-6} alkyloxycarbonyl, C_{1-6} alkylcarbonyl, formyl, $-NR^{13}R^{14}$ or R^7 ;
- R^5 is hydrogen; aryl; formyl; C_{1-6} alkylcarbonyl; C_{1-6} alkyl; C_{1-6} alkyloxycarbonyl; C_{1-6} alkyl substituted with formyl, C_{1-6} alkylcarbonyl, C_{1-6} alkyloxycarbonyl or C_{1-6} alkylcarbonyloxy; C_{1-6} alkyloxy C_{1-6} alkylcarbonyl substituted with C_{1-6} alkyloxycarbonyl;
- R^6 is C_{1-4} alkyl, $NR^{13}R^{14}$ or polyhalo C_{1-4} alkyl;
- R^7 is a monocyclic, bicyclic or tricyclic saturated, partially saturated or aromatic carbocycle or a monocyclic, bicyclic or tricyclic saturated, partially saturated or aromatic heterocycle, wherein each of said carbocyclic or heterocyclic ring systems may optionally be substituted with one, two, three, four or five substituents each independently selected from halo, hydroxy, mercapto, C_{1-6} alkyl, hydroxy C_{1-6} alkyl, amino C_{1-6} alkyl, mono or di(C_{1-6} alkyl)amino C_{1-6} alkyl, formyl, C_{1-6} alkylcarbonyl, C_{3-7} cycloalkyl, C_{1-6} alkyloxy, C_{1-6} alkyloxycarbonyl, C_{1-6} alkylthio, cyano, nitro, polyhalo C_{1-6} alkyl, polyhalo C_{1-6} alkyloxy, aminocarbonyl, $-CH(=N-O-R^8)$, R^{7a} , $-X_3-R^{7a}$ or $R^{7a}-C_{1-4}$ alkyl;
- R^{7a} is a monocyclic, bicyclic or tricyclic saturated, partially saturated or aromatic carbocycle or a monocyclic, bicyclic or tricyclic saturated, partially saturated or aromatic heterocycle, wherein each of said carbocyclic or heterocyclic ring systems may optionally be substituted with one, two, three, four or five substituents each independently selected from halo, hydroxy, mercapto, C_{1-6} alkyl, hydroxy C_{1-6} alkyl, amino C_{1-6} alkyl, mono or di(C_{1-6} alkyl)amino C_{1-6} alkyl, formyl, C_{1-6} alkylcarbonyl, C_{3-7} cycloalkyl, C_{1-6} alkyloxy, C_{1-6} alkyloxycarbonyl, C_{1-6} alkylthio, cyano, nitro, polyhalo C_{1-6} alkyl, polyhalo C_{1-6} alkyloxy, $-C(=O)-NR^{13}R^{14}$, $-CH(=N-O-R^8)$;
- R^8 is hydrogen, C_{1-4} alkyl, aryl or aryl C_{1-4} alkyl;
- R^9 and R^{10} each independently are hydrogen; hydroxy; C_{1-6} alkyl; C_{1-6} alkyloxy; C_{1-6} alkylcarbonyl; C_{1-6} alkyloxycarbonyl; $NR^{13}R^{14}$; $-C(=O)-NR^{13}R^{14}$; $-CH(=NR^{11})$ or R^7 , wherein each of the aforementioned C_{1-6} alkyl groups may optionally and each individually be substituted with one or two substituents each independently selected from hydroxy, C_{1-6} alkyloxy, hydroxy C_{1-6} alkyloxy, carboxyl, C_{1-6} alkyloxycarbonyl, cyano, imino, $NR^{13}R^{14}$, polyhalomethyl, polyhalomethyloxy, polyhalomethylthio, $-S(=O)_pR^6$, $-NH-S(=O)_pR^6$, $-C(=O)R^6$, $-NHC(=O)H$, $-C(=O)NHNH_2$, $-NHC(=O)R^6$, $-C(=NH)R^6$, R^7 ; or
- R^9 and R^{10} may be taken together to form a bivalent or trivalent radical of formula
- $-CH_2-CH_2-CH_2-CH_2-$ (d-1)
 $-CH_2-CH_2-CH_2-CH_2-CH_2-$ (d-2)
 $-CH_2-CH_2-O-CH_2-CH_2-$ (d-3)

-4-



- 5 R^{11} is cyano; C_{1-4} alkylcarbonyl; C_{1-4} alkyloxycarbonyl; $-\text{C}(=\text{O})-\text{NR}^{13}\text{R}^{14}$; or C_{1-4} alkyl optionally substituted with C_{1-4} alkyloxy, cyano, $\text{NR}^{13}\text{R}^{14}$ or $-\text{C}(=\text{O})-\text{NR}^{13}\text{R}^{14}$; R^{12} is hydrogen or C_{1-4} alkyl; R^{13} and R^{14} each independently are hydrogen, Het, C_{1-6} alkyl optionally substituted with cyano or aminocarbonyl, C_{2-6} alkenyl optionally substituted with cyano or aminocarbonyl, C_{2-6} alkynyl optionally substituted with cyano or aminocarbonyl;
- 10 R^{15} is C_{1-6} alkyl optionally substituted with cyano or $-\text{C}(=\text{O})-\text{NR}^{13}\text{R}^{14}$; R^{16} is R^7 or C_{1-6} alkyl optionally substituted with cyano or $-\text{C}(=\text{O})-\text{NR}^{13}\text{R}^{14}$; R^{17} , if present, each independently is cyano, halo, hydroxy, $-\text{C}(=\text{O})-\text{NR}^{13}\text{R}^{14}$, C_{1-6} alkyl optionally substituted with one or more substituents independently selected from cyano, $-\text{C}(=\text{O})-\text{NR}^{13}\text{R}^{14}$ or halo; C_{2-6} alkenyl optionally substituted with one or more substituents independently selected from cyano, $-\text{C}(=\text{O})-\text{NR}^{13}\text{R}^{14}$ or halo; C_{2-6} alkynyl optionally substituted with one or more substituents independently selected from cyano, $-\text{C}(=\text{O})-\text{NR}^{13}\text{R}^{14}$ or halo; and, where possible, R^{17} may also be attached to the $-\text{b}^1-\text{b}^2-\text{b}^3-$ moiety by a double bond whereby R^{17} is then $=\text{O}$, $=\text{S}$, $=\text{NH}$, $=\text{N}-\text{R}^{15}$, $=\text{N}-\text{R}^7$, $=\text{N}-\text{O}-\text{R}^{15}$, $=\text{N}-\text{O}-\text{R}^7$, $=\text{CH}_2$, $=\text{CH}-\text{C}(=\text{O})-\text{NR}^{13}\text{R}^{14}$, $=\text{CH}-\text{R}^7$, or $=\text{CH}-\text{R}^{15}$; wherein $=\text{CH}_2$ may optionally be substituted with cyano, hydroxy, halo, nitro;
- 15 Q represents hydrogen, C_{1-6} alkyl, halo, polyhalo C_{1-6} alkyl, $-\text{C}(=\text{O})-\text{NR}^{13}\text{R}^{14}$, or $-\text{NR}^9\text{R}^{10}$; Z is C-Y, wherein, Y represents hydrogen, hydroxy, halo, C_{1-6} alkyl, C_{3-7} cycloalkyl, C_{1-6} alkyloxy, C_{1-6} alkyloxycarbonyl, carbonyl, cyano, nitro, $\text{NR}^{13}\text{R}^{14}$, polyhalomethyl, polyhalomethyloxy, polyhalomethylthio, $-\text{S}(=\text{O})_p\text{R}^8$, $-\text{NH}-\text{S}(=\text{O})\text{R}^8$, $-\text{NH}-\text{SO}_2-\text{R}^8$, $-\text{NH}-\text{SO}_2-(\text{C}_{1-4}\text{alkanediyl})-\text{CO}-\text{N}(\text{R}^8)_2$, $-\text{C}(=\text{O})\text{R}^8$, $-\text{NHC}(=\text{O})\text{H}$, $-\text{C}(=\text{O})\text{NHNH}_2$, $-\text{NHC}(=\text{O})\text{R}^8$, $-\text{C}(=\text{O})-\text{NH}-\text{R}^8$, $-\text{C}(=\text{NH})\text{R}^8$, aryl or C_{2-6} alkenyl optionally substituted with one or more halo atoms;
- 20 C_{2-6} alkynyl optionally substituted with one or more halo atoms; C_{1-6} alkyl substituted with cyano, or with $-\text{C}(=\text{O})\text{R}^8$, aryl is phenyl or phenyl substituted with one, two, three, four or five substituents each independently selected from halo, hydroxy, mercapto, C_{1-6} alkyl, hydroxy C_{1-6} alkyl, C_{1-6} alkyl $\text{NR}^{13}\text{R}^{14}$, C_{1-6} alkylcarbonyl, C_{3-7} cycloalkyl, C_{1-6} alkyloxy, C_{1-6} alkyloxycarbonyl, C_{1-6} alkylthio, cyano, nitro, polyhalo C_{1-6} alkyl, polyhalo C_{1-6} alkyloxy, $-\text{C}(=\text{O})-\text{NR}^{13}\text{R}^{14}$, R^7 or $-\text{X}_3-\text{R}^7$;
- 25 Het is a monocyclic, bicyclic or tricyclic saturated, partially saturated or aromatic heterocycle, wherein each of said carbocyclic or heterocyclic ring systems may

-5-

optionally be substituted with one, two, three, four or five substituents each independently selected from halo, hydroxy, mercapto, C₁₋₆alkyl, hydroxyC₁₋₆alkyl, aminoC₁₋₆alkyl, mono or di(C₁₋₆alkyl)aminoC₁₋₆alkyl, formyl, C₁₋₆alkylcarbonyl, C₃₋₇cycloalkyl, C₁₋₆alkyloxy, C₁₋₆alkyloxycarbonyl, C₁₋₆alkylthio, cyano, nitro,
 5 polyhaloC₁₋₆alkyl, polyhaloC₁₋₆alkyloxy, -C(=O)-NR¹³R¹⁴, -CH(=N-O-R⁸).

As used hereinbefore or hereinafter C₁₋₄alkyl as a group or part of a group defines straight or branched chain saturated hydrocarbon radicals having from 1 to 4 carbon atoms such as methyl, ethyl, propyl, 1-methylethyl, butyl; C₁₋₆alkyl as a group or part of
 10 a group defines straight or branched chain saturated hydrocarbon radicals having from 1 to 6 carbon atoms such as the group defined for C₁₋₄alkyl and pentyl, hexyl, 2-methylbutyl and the like; C₂₋₆alkyl as a group or part of a group defines straight or branched chain saturated hydrocarbon radicals having from 2 to 6 carbon atoms such as ethyl, propyl, 1-methylethyl, butyl, pentyl, hexyl, 2-methylbutyl and the like; C₁₋₄alkanediyl
 15 defines straight or branched chain saturated bivalent hydrocarbon radicals having from 1 to 4 carbon atoms such as methylene, 1,2-ethanediyl or 1,2-ethylidene, 1,3-propanediyl or 1,3-propylidene, 1,4-butanediyl or 1,4-butyldiene and the like; C₃₋₇cycloalkyl is generic to cyclopropyl, cyclobutyl, cyclopentyl, cyclohexyl and cycloheptyl; C₂₋₆alkenyl defines straight and branched chain hydrocarbon radicals having from 2 to 6 carbon
 20 atoms containing a double bond such as ethenyl, propenyl, butenyl, pentenyl, hexenyl and the like; C₂₋₆alkynyl defines straight and branched chain hydrocarbon radicals having from 2 to 6 carbon atoms containing a triple bond such as ethynyl, propynyl, butynyl, pentynyl, hexynyl and the like.

25 In a number of instances the radicals C₁₋₆alkynyl, C₂₋₆alkenyl, or C₂₋₆alkynyl may be substituted with one or more substituents. In that instance there can be 1, 2, 3, 4, 5, 6 and more substituents, the number in some cases being limited by the number of carbon atoms and the degree of unsaturation of the hydrocarbon radical. Preferably, the radicals C₁₋₆alkynyl, C₂₋₆alkenyl, or C₂₋₆alkynyl are substituted with up to 3 substituents.

30 A monocyclic, bicyclic or tricyclic saturated carbocycle represents a ring system consisting of 1, 2 or 3 rings, said ring system being composed of only carbon atoms and said ring system containing only single bonds; a monocyclic, bicyclic or tricyclic partially saturated carbocycle represents a ring system consisting of 1, 2 or 3 rings, said ring
 35 system being composed of only carbon atoms and comprising at least one double bond provided that the ring system is not an aromatic ring system; a monocyclic, bicyclic or tricyclic aromatic carbocycle represents an aromatic ring system consisting of 1, 2 or 3 rings, said ring system being composed of only carbon atoms; the term aromatic is well

-6-

known to a person skilled in the art and designates cyclically conjugated systems of $4n + 2$ electrons, that is with 6, 10, 14 etc. π -electrons (rule of Hückel) ; a monocyclic, bicyclic or tricyclic saturated heterocycle represents a ring system consisting of 1, 2 or 3 rings and comprising at least one heteroatom selected from O, N or S, said ring system
5 containing only single bonds; a monocyclic, bicyclic or tricyclic partially saturated heterocycle represents a ring system consisting of 1, 2 or 3 rings and comprising at least one heteroatom selected from O, N or S, and at least one double bond provided that the ring system is not an aromatic ring system; a monocyclic, bicyclic or tricyclic aromatic heterocycle represents an aromatic ring system consisting of 1, 2 or 3 rings and
10 comprising at least one heteroatom selected from O, N or S.

Particular examples of monocyclic, bicyclic or tricyclic saturated carbocycles are cyclopropyl, cyclobutyl, cyclopentyl, cyclohexyl, cycloheptyl, cyclooctyl, bicyclo[4,2,0]-octanyl, cyclononanyl, cyclodecanyl, decahydronaphthalenyl, tetradecahydroanthracenyl
15 and the like. Preferred are cyclopropyl, cyclobutyl, cyclopentyl, cyclohexyl, cycloheptyl; more preferred are cyclopentyl, cyclohexyl, cycloheptyl.

Particular examples of monocyclic, bicyclic or tricyclic partially saturated carbocycles are cyclopropenyl, cyclobutenyl, cyclopentenyl, cyclohexenyl, cycloheptenyl, cyclooctenyl, bicyclo[4,2,0]octenyl, cyclononenyl, cyclodecenyl, octahydronaphthalenyl, 1,2,3,4-tetrahydronaphthalenyl, 1,2,3,4,4a,9,9a,10-octahydro-anthracenyl and the like.
20 Preferred are cyclopentenyl, cyclohexenyl, cycloheptenyl.

Particular examples of monocyclic, bicyclic or tricyclic aromatic carbocycles are phenyl, naphthalenyl, anthracenyl. Preferred is phenyl.
25

Particular examples of monocyclic, bicyclic or tricyclic saturated heterocycles are tetrahydrofuranyl, pyrrolidinyl, dioxolanyl, imidazolidinyl, thiazolidinyl, tetrahydrothienyl, dihydrooxazolyl, isothiazolidinyl, isoxazolidinyl, oxadiazolidinyl, triazolidinyl, thiadiazolidinyl, pyrazolidinyl, piperidinyl, hexahydropyrimidinyl, hexahydropyrazinyl, dioxanyl, morpholinyl, dithianyl, thiomorpholinyl, piperazinyl, trithianyl, decahydroquinolinyl, octahydroindolyl and the like. Preferred are tetrahydrofuranyl, pyrrolidinyl, dioxolanyl, imidazolidinyl, thiazolidinyl, dihydrooxazolyl, triazolidinyl, piperidinyl, dioxanyl, morpholinyl, thiomorpholinyl, piperazinyl.
30 Particularly preferred are tetrahydrofuranyl, pyrrolidinyl, dioxolanyl, piperidinyl, dioxanyl, morpholinyl, thiomorpholinyl, piperazinyl.
35

Particular examples of monocyclic, bicyclic or tricyclic partially saturated heterocycles are pyrrolinyl, imidazolinyl, pyrazolinyl, 2,3-dihydrobenzofuranyl, 1,3-benzodioxolyl,

-7-

2,3-dihydro-1,4-benzodioxinyl, indolinyl and the like. Preferred are pyrrolinyl, imidazolinyl, 2,3-dihydrobenzofuranyl, 1,3-benzodioxolyl, indolinyl.

- Particular examples of monocyclic, bicyclic or tricyclic aromatic heterocycles are azetyl,
- 5 oxetylidenyl, pyrrolyl, furyl, thienyl, imidazolyl, oxazolyl, isoxazolyl, thiazolyl, isothiazolyl, pyrazolyl, triazolyl, thiadiazolyl, oxadiazolyl, tetrazolyl, pyridyl, pyrimidinyl, pyrazinyl, pyridazinyl, triazinyl, pyranyl, benzofuryl, isobenzofuryl, benzo-
- thienyl, isobenzothienyl, indoliziny, indolyl, isoindolyl, benzoxazolyl, benzimidazolyl, indazolyl, benzisoxazolyl, benzisothiazolyl, benzopyrazolyl, benzoxadiazolyl,
- 10 benzothiadiaazolyl, benzotriazolyl, purinyl, quinoliny, isoquinoliny, cinnoliny, quinoliziny, phthalaziny, quinoxaliny, quinazoliny, naphthiridiny, pteridiny, benzopyrany, pyrrolopyridyl, thienopyridyl, furopyridyl, isothiazolopyridyl, thiazolopyridyl, isoxazolopyridyl, oxazolopyridyl, pyrazolopyridyl, imidazopyridyl, pyrrolopyraziny, thienopyraziny, furopyraziny, isothiazolopyraziny, thiazolopyraziny,
- 15 isoxazolopyraziny, oxazolopyraziny, pyrazolopyraziny, imidazopyraziny, pyrrolopyrimidinyl, thienopyrimidinyl, fuopyrimidinyl, isothiazolopyrimidinyl, thiazolopyrimidinyl, isoxazolopyrimidinyl, oxazolopyrimidinyl, pyrazolopyrimidinyl, imidazopyrimidinyl, pyrrolopyridaziny, thienopyridaziny, fuopyridaziny, isothiazolopyridaziny, thiazolopyridaziny, isoxazolopyridaziny, oxazolopyridaziny,
- 20 pyrazolopyridaziny, imidazopyridaziny, oxadiazolopyridyl, thiadiazolopyridyl, triazolopyridyl, oxadiazolopyraziny, thiadiazolopyraziny, triazolopyraziny, oxadiazolopyrimidinyl, thiadiazolopyrimidinyl, triazolopyrimidinyl, oxadiazolo-
- pyridaziny, thiadiazolopyridaziny, triazolopyridaziny, imidazooxazolyl, imidazothiazolyl, imidazoimidazolyl, isoxazolotriazinyl, isothiazolotriazinyl,
- 25 pyrazolotriazinyl, oxazolotriazinyl, thiazolotriazinyl, imidazotriazinyl, oxadiazolotriazinyl, thiadiazolotriazinyl, triazolotriazinyl, carbazolyl, acridiny, phenaziny, phenothiaziny, phenoxaziny and the like.

Preferred aromatic heterocycles are monocyclic or bicyclic aromatic heterocycles.

- 30 Interesting monocyclic, bicyclic or tricyclic aromatic heterocycles are pyrrolyl, furyl, thienyl, imidazolyl, oxazolyl, isoxazolyl, thiazolyl, isothiazolyl, pyrazolyl, triazolyl, thiadiazolyl, oxadiazolyl, tetrazolyl, pyridyl, pyrimidinyl, pyraziny, pyridazinyl, triazinyl, pyranyl, benzofuryl, isobenzofuryl, benzo-
- thienyl, isobenzothienyl, indolyl, isoindolyl, benzoxazolyl, benzimidazolyl, indazolyl, benzisoxazolyl, benzisothiazolyl,
- 35 benzopyrazolyl, benzoxadiazolyl, benzothiadiaazolyl, benzotriazolyl, purinyl, quinoliny, isoquinoliny, phthalaziny, quinoxaliny, quinazoliny, benzopyrany, pyrrolopyridyl, thienopyridyl, furopyridyl, isothiazolopyridyl, thiazolopyridyl, isoxazolopyridyl, oxazolopyridyl, pyrazolopyridyl, imidazopyridyl, pyrrolopyraziny, thienopyraziny,

5 fuoropyrazinyl, isothiazolopyrazinyl, thiazolopyrazinyl, isoxazolopyrazinyl, oxazolo-
pyrazinyl, pyrazolopyrazinyl, imidazopyrazinyl, pyrrolopyrimidinyl, thienopyrimidinyl,
fuoropyrimidinyl, isothiazolopyrimidinyl, thiazolopyrimidinyl, isoxazolopyrimidinyl,
oxazolopyrimidinyl, pyrazolopyrimidinyl, imidazopyrimidinyl, oxadiazolopyridyl,
10 thiadiazolopyridyl, triazolopyridyl, oxadiazolopyrazinyl, thiadiazolopyrazinyl,
triazolopyrazinyl, oxadiazolopyrimidinyl, thiadiazolopyrimidinyl, triazolopyrimidinyl,
carbazolyl, acridinyl, phenothiazinyl, phenoxazinyl and the like.

10 Particularly interesting aromatic heterocycles are pyrrolyl, furyl, thienyl, imidazolyl,
oxazolyl, isoxazolyl, thiazolyl, isothiazolyl, pyrazolyl, triazolyl, thiadiazolyl, oxadiazolyl,
tetrazolyl, pyridyl, pyrimidinyl, pyrazinyl, pyridazinyl, triazinyl, pyranyl, benzofuryl,
isobenzofuryl, benzothienyl, isobenzothienyl, indolyl, isoindolyl, benzoxazolyl,
benzimidazolyl, indazolyl, benzisoxazolyl, benzisothiazolyl, benzopyrazolyl,
15 benzoxadiazolyl, benzothiadiazolyl, benzotriazolyl, purinyl, quinolinyl, isoquinolinyl,
phthalazinyl, quinoxalinyl, quinazolinyl, and the like.

20 As used herein before, the term (=O) forms a carbonyl moiety when attached to a
carbon atom, a sulfoxide moiety when attached to a sulfur atom and a sulfonyl moiety
when two of said terms are attached to a sulfur atom.

25 The term halo is generic to fluoro, chloro, bromo and iodo. As used in the foregoing and
hereinafter, polyhalomethyl as a group or part of a group is defined as mono- or
polyhalosubstituted methyl, in particular methyl with one or more fluoro atoms, for
example, difluoromethyl or trifluoromethyl; polyhaloC₁₋₄alkyl or polyhaloC₁₋₆alkyl as a
group or part of a group is defined as mono- or polyhalosubstituted C₁₋₄alkyl or
C₁₋₆alkyl, for example, the groups defined in halomethyl, 1,1-difluoro-ethyl and the like.
In case more than one halogen atoms are attached to an alkyl group within the definition
of polyhalomethyl, polyhaloC₁₋₄alkyl or polyhaloC₁₋₆alkyl, they may be the same or
different.

30 The term heterocycle in the definition of R⁷ or R^{7a} is meant to include all the possible
isomeric forms of the heterocycles, for instance, pyrrolyl comprises 1*H*-pyrrolyl and 2*H*-
pyrrolyl.

35 The carbocycle or heterocycle in the definition of R⁷ or R^{7a} may be attached to the
remainder of the molecule of formula (I) through any ring carbon or heteroatom as
appropriate, if not otherwise specified. Thus, for example, when the heterocycle is
imidazolyl, it may be 1-imidazolyl, 2-imidazolyl, 4-imidazolyl and the like, or when the
carbocycle is naphthalenyl, it may be 1-naphthalenyl, 2-naphthalenyl and the like.

When any variable (eg. R^7 , heteroatom, X_2) occurs more than one time in any constituent, each definition is independent.

- 5 Lines drawn from substituents into ring systems indicate that the bond may be attached to any of the suitable ring atoms.

For therapeutic use, salts of the compounds of formula (I) are those wherein the counterion is pharmaceutically acceptable. However, salts of acids and bases which are
10 non-pharmaceutically acceptable may also find use, for example, in the preparation or purification of a pharmaceutically acceptable compound. All salts, whether pharmaceutically acceptable or not are included within the ambit of the present invention.

- 15 The pharmaceutically acceptable addition salts as mentioned hereinabove are meant to comprise the therapeutically active non-toxic acid addition salt forms, which the compounds of formula (I) are able to form. The latter can conveniently be obtained by treating the base form with such appropriate acids as inorganic acids, for example, hydrohalic acids, e.g. hydrochloric, hydrobromic and the like; sulfuric acid; nitric acid;
20 phosphoric acid and the like; or organic acids, for example, acetic, propanoic, hydroxyacetic, 2-hydroxypropanoic, 2-oxopropanoic, oxalic, malonic, succinic, maleic, fumaric, malic, tartaric, 2-hydroxy-1,2,3-propanetricarboxylic, methanesulfonic, ethanesulfonic, benzenesulfonic, 4-methylbenzenesulfonic, cyclohexanesulfamic, 2-hydroxybenzoic, 4-amino-2-hydroxybenzoic and the like acids. Conversely the salt form can be converted
25 by treatment with alkali into the free base form.

- The compounds of formula (I) containing acidic protons may be converted into their therapeutically active non-toxic metal or amine addition salt forms by treatment with appropriate organic and inorganic bases. Appropriate base salt forms comprise, for
30 example, the ammonium salts, the alkali and earth alkaline metal salts, e.g. the lithium, sodium, potassium, magnesium, calcium salts and the like, salts with organic bases, e.g. primary, secondary and tertiary aliphatic and aromatic amines such as methylamine, ethylamine, propylamine, isopropylamine, the four butylamine isomers, dimethylamine, diethylamine, diethanolamine, dipropylamine, diisopropylamine, di-n-butylamine,
35 pyrrolidine, piperidine, morpholine, trimethylamine, triethylamine, tripropylamine, quinuclidine, pyridine, quinoline and isoquinoline, the benzathine, *N*-methyl-D-glucamine, 2-amino-2-(hydroxymethyl)-1,3-propanediol, hydrabamine salts, and salts with amino acids such as, for example, arginine, lysine and the like. Conversely the salt form can be converted by treatment with acid into the free acid form.

-10-

The term addition salt also comprises the hydrates and solvent addition forms (solvates) which the compounds of formula (I) are able to form. Examples of such forms are e.g. hydrates, alcoholates and the like.

- 5 The term "quaternary amine" as used hereinbefore defines the quaternary ammonium salts which the compounds of formula (I) are able to form by reaction between a basic nitrogen of a compound of formula (I) and an appropriate quaternizing agent, such as, for example, an optionally substituted alkylhalide, arylhalide or arylalkylhalide, e.g. methyl iodide or benzyl iodide. Other reactants with good leaving groups may also be used, such as alkyl trifluoromethanesulfonates, alkyl methanesulfonates, and alkyl p-toluenesulfonates. A quaternary amine has a positively charged nitrogen. Pharmaceutically acceptable counterions include chloro, bromo, iodo, trifluoroacetate and acetate. The counterion of choice can be introduced using ion exchange resins.
- 10
- 15 The *N*-oxide forms of the present compounds are meant to comprise the compounds of formula (I) wherein one or several tertiary nitrogen atoms are oxidized to the so-called *N*-oxide.

- 20 It will be appreciated that some of the compounds of formula (I) and their *N*-oxides, addition salts, quaternary amines and stereochemically isomeric forms may contain one or more centers of chirality and exist as stereochemically isomeric forms.

- The term "stereochemically isomeric forms" as used hereinbefore defines all the possible stereoisomeric forms which the compounds of formula (I), and their *N*-oxides, addition salts, quaternary amines or physiologically functional derivatives may possess. Unless otherwise mentioned or indicated, the chemical designation of compounds denotes the mixture of all possible stereochemically isomeric forms, said mixtures containing all diastereomers and enantiomers of the basic molecular structure as well as each of the individual isomeric forms of formula (I) and their *N*-oxides, salts, solvates or quaternary amines substantially free, *i.e.* associated with less than 10%, preferably less than 5%, in particular less than 2% and most preferably less than 1% of the other isomers. Thus, when a compound of formula (I) is for instance specified as (E), this means that the compound is substantially free of the (Z) isomer.
- 25
- 30

- 35 In particular, stereogenic centers may have the R- or S-configuration; substituents on bivalent cyclic (partially) saturated radicals may have either the *cis*- or *trans*-configuration. Compounds encompassing double bonds can have an E (entgegen) or Z (zusammen)-stereochemistry at said double bond. The terms *cis*, *trans*, R, S, E and Z are well known to a person skilled in the art.

-11-

Stereochemically isomeric forms of the compounds of formula (I) obviously are intended to be embraced within the scope of this invention.

- 5 For some of the compounds of formula (I), their prodrugs, *N*-oxides, salts, solvates, quaternary amines or metal complexes and the intermediates used in the preparation thereof, the absolute stereochemical configuration was not experimentally determined. In these cases the stereoisomeric form, which was first isolated is designated as "A" and the second as "B", without further reference to the actual stereochemical configuration.
- 10 However, said "A" and "B" stereoisomeric forms can be unambiguously characterized by for instance their optical rotation in case "A" and "B" have an enantiomeric relationship. A person skilled in the art is able to determine the absolute configuration of such compounds using art-known methods such as, for example, X-ray diffraction. In case "A" and "B" are stereoisomeric mixtures, they can be further separated whereby
- 15 the respective first fractions isolated are designated "A1" and "B1" and the second as "A2" and "B2", without further reference to the actual stereochemical configuration.

- Some of the compounds of formula (I) may also exist in their tautomeric form. Such forms although not explicitly indicated in the above formula are intended to be included
- 20 within the scope of the present invention.

- Whenever used hereinafter, the term "compounds of formula (I)" is meant to also include their *N*-oxide forms, their salts, their quaternary amines and their stereochemically isomeric forms. Of special interest are those compounds of formula (I)
- 25 which are stereochemically pure.


- Whenever used hereinbefore or hereinafter that substituents can be selected each independently out of a list of numerous definitions, such as for example for R^9 and R^{10} , all possible combinations are intended which are chemically possible and which lead to
- 30 chemically stable molecules.

Subgroups of the compounds of formula (I) that are of interest are those wherein one or more of the following limitations (a) - (v) apply.

- (a) $-a^1=a^2-a^3=a^4-$ represents a bivalent radical of formula
- 35 $-CH=CH-CH=CH-$ (a-1);
- (b) n is 0, 1, 2, 3;
- (c) m is 0, 1 or 2;
- (d) R^1 is hydrogen; formyl; C_{1-6} alkylcarbonyl; C_{1-6} alkyl; C_{1-6} alkyloxy carbonyl; C_{1-6} alkylcarbonyl, C_{1-6} alkyloxy carbonyl;

-12-

- (e) each R^2 independently is hydroxy, halo, C_{1-6} alkyl optionally substituted with cyano or with $-C(=O)R^6$, C_{3-7} cycloalkyl, C_{2-6} alkenyl optionally substituted with one or more halogen atoms or cyano, C_{2-6} alkynyl optionally substituted with one or more halogen atoms or cyano, C_{1-6} alkyloxycarbonyl, carboxyl, cyano, nitro, amino, mono(C_{1-6} alkyl)amino, di(C_{1-6} alkyl)amino, polyhalomethyl, polyhalomethylthio, $-S(=O)_pR^6$, $-NH-S(=O)_pR^6$, $-C(=O)R^6$, $-NHC(=O)H$, $-C(=O)NHNH_2$, $-NHC(=O)R^6$, $-C(=NH)R^6$ or a radical of formula



- wherein each A_1 independently is N, CH or CR^6 ; and
 A_2 is NH, O, S or NR^6 ;
- (f) X_1 is $-NR^5$ -, $-NH-NH$ -, $-N=N$ -, $-O$ -, $-C(=O)$ -, C_{1-4} alkanediyl, $-CHOH$ -, $-S$ -, $-S(=O)_p$ -, $-NR^{13}-C(=O)$ -, $-C(=O)-NR^{13}$ -, $-X_2-C_{1-4}$ alkanediyl- or $-C_{1-4}$ alkanediyl- X_2 ;
- (g) X_2 is $-NR^5$ -, $-O$ -;
- (h) R^3 is hydrogen, halo, C_{1-6} alkyl, $NR^{13}R^{14}$, $-C(=O)-NR^{13}R^{14}$, $-C(=O)-R^{15}$, $-X_3-R^7$; C_{1-6} alkyl substituted with one or more substituents each independently selected from cyano, R^7 or $-C(=O)-NR^9R^{10}$; C_{2-6} alkenyl substituted with one or more substituents each independently selected from halo, cyano or $-C(=O)-NR^9R^{10}$ or R^7 ; or C_{2-6} alkynyl substituted with one or more substituents each independently selected from halo, cyano, $-C(=O)-NR^9R^{10}$ or R^7 ;
- (i) X_3 is $-NR^5$ -, $-NH-NH$ -, $-N=N$ -, $-O$ - or $-S$ -;
- (j) R^4 is halo, hydroxy, C_{1-6} alkyl, C_{2-6} alkenyl, C_{2-6} alkynyl, C_{1-6} alkyloxy, cyano, nitro, polyhalo C_{1-6} alkyl, polyhalo C_{1-6} alkyloxy, $-C(=O)-NR^{13}R^{14}$, C_{1-6} alkyloxycarbonyl, C_{1-6} alkylcarbonyl, formyl, $-NR^{13}R^{14}$ or R^7 ;
- (k) R^5 is hydrogen; formyl; C_{1-6} alkylcarbonyl; C_{1-6} alkyl or C_{1-6} alkyloxycarbonyl;
- (l) R^6 is C_{1-4} alkyl, $NR^{13}R^{14}$ or polyhalo C_{1-4} alkyl;
- (m) R^7 is a monocyclic or bicyclic, partially saturated or aromatic carbocycle or a monocyclic or bicyclic, partially saturated or aromatic heterocycle, wherein each of said carbocyclic or heterocyclic ring systems may optionally be substituted with one, two or three substituents each independently selected from halo, hydroxy, mercapto, C_{1-6} alkyl, hydroxy C_{1-6} alkyl, amino C_{1-6} alkyl, C_{1-6} alkylcarbonyl, C_{1-6} alkyloxy, C_{1-6} alkyloxycarbonyl, C_{1-6} alkylthio, cyano, nitro, polyhalo C_{1-6} alkyl, polyhalo C_{1-6} alkyloxy or aminocarbonyl;
- (n) R^8 is hydrogen, C_{1-4} alkyl or aryl C_{1-4} alkyl;
- (o) R^9 and R^{10} each independently are hydrogen; C_{1-6} alkyl; C_{1-6} alkyloxy; C_{1-6} alkylcarbonyl or C_{1-6} alkyloxycarbonyl;
- (p) R^{13} and R^{14} each independently are hydrogen or C_{1-6} alkyl;

-13-

- (q) R^{15} is C_{1-6} alkyl optionally substituted with cyano or $-C(=O)-NR^{13}R^{14}$;
- (r) R^{17} is cyano, halo, hydroxy, $-C(=O)-NR^{13}R^{14}$, C_{1-6} alkyl optionally substituted with cyano, $-C(=O)-NR^{13}R^{14}$ or halo; C_{2-6} alkenyl optionally substituted with cyano or $-C(=O)-NR^{13}R^{14}$; C_{2-6} alkynyl optionally substituted with cyano or $-C(=O)-NR^{13}R^{14}$;
- 5 and, where possible, R^{17} may also be attached to the $-b^1-b^2-b^3-$ moiety by a double bond whereby R^{17} is then $=O$, $=S$, $=NH$, $=N-R^{15}$, $=N-R^7$, $=N-O-R^{15}$, $=N-O-R^7$, $=CH_2$, $=CH-C(=O)-NR^{13}R^{14}$, $=CH-R^7$, or $=CH-R^{15}$; wherein $=CH_2$ may optionally be substituted with cyano, hydroxy, halo, nitro;
- (s) Q represents hydrogen, C_{1-6} alkyl or $-NR^9R^{10}$;
- 10 (t) Y represents hydrogen, hydroxy, halo, C_{1-6} alkyl, C_{1-6} alkyloxy, cyano, nitro, $NR^{13}R^{14}$, polyhalomethyloxy, $-NH-SO_2-R^8$, $-NH-SO_2-(C_{1-4}alkanedyl)-CO-N(R^8)_2$; or Y is C_{1-6} alkyl substituted with cyano or with $-C(=O)R^8$;
- (u) aryl is phenyl or phenyl substituted with one, two or three substituents each independently selected from halo, hydroxy, mercapto, C_{1-6} alkyl, hydroxy C_{1-6} alkyl,
- 15 C_{1-6} alkyl $NR^{13}R^{14}$, C_{1-6} alkylcarbonyl, C_{1-6} alkyloxy, C_{1-6} alkyloxycarbonyl, C_{1-6} alkylthio, cyano, nitro, polyhalo C_{1-6} alkyl, polyhalo C_{1-6} alkyloxy, $-C(=O)-NR^{13}R^{14}$, R^7 or $-X_3-R^7$;
- (v) Het is a monocyclic or bicyclic, partially saturated or aromatic heterocycle, wherein each of said carbocyclic or heterocyclic ring systems may optionally be substituted
- 20 with one, two or three substituents each independently selected from halo, hydroxy, mercapto, C_{1-6} alkyl, hydroxy C_{1-6} alkyl, amino C_{1-6} alkyl, C_{1-6} alkylcarbonyl, C_{1-6} alkyloxy, C_{1-6} alkyloxycarbonyl, C_{1-6} alkylthio, cyano, nitro, polyhalo C_{1-6} alkyl, polyhalo C_{1-6} alkyloxy.
- 25 A particularly interesting subgroup of compounds of formula (I) are those wherein all of the above limitations (a) - (v) apply.

Of further interest are subgroups of the compounds of formula (I) wherein one or more of the afore mentioned limitations (a) - (v) optionally apply and one or more of the

30 following limitations (a') - (v') apply :

(a') $-a^1=a^2-a^3=a^4-$ represents a bivalent radical of formula

$$-CH=CH-CH=CH- \quad (a-1);$$

(b') n is 1 or 2;


35 (c') m is 1 or 2;

(d') R^1 is hydrogen; C_{1-6} alkyl;

(e') each R^2 independently is hydroxy, halo, C_{1-6} alkyl optionally substituted with cyano or with $-C(=O)R^6$, C_{2-6} alkenyl optionally substituted with cyano, C_{2-6} alkynyl optionally substituted with cyano, C_{1-6} alkyloxycarbonyl, carboxyl, cyano, nitro,

-14-

amino, mono(C₁₋₆alkyl)amino, di(C₁₋₆alkyl)amino, -S(=O)_pR⁶, -NH-S(=O)_pR⁶, -C(=O)R⁶, -NHC(=O)H, -C(=O)NHNH₂, -NHC(=O)R⁶, -C(=NH)R⁶ or a radical of formula



- 5 wherein each A₁ independently is N, CH or CR⁶; and no more than two A₁ are N;
A₂ is NH, O, S or NR⁶;
- (f') X₁ is -NR⁵-, -NH-NH-, -N=N-, -O-, -C(=O)-, C₁₋₄alkanediyl, -CHOH-,
-NR¹³-C(=O)-, -C(=O)-NR¹³-, -X₂-C₁₋₄alkanediyl- or -C₁₋₄alkanediyl-X₂-;
- (g') X₂ is -NR⁵-, -O-;
- 10 (h') R³ is hydrogen, halo, C₁₋₆alkyl, NR¹³R¹⁴, -C(=O)-NR¹³R¹⁴, -C(=O)-R¹⁵, -X₃-R⁷;
C₁₋₆alkyl substituted with one or two substituents each independently selected from
cyano, R⁷ or -C(=O)-NR⁹R¹⁰; C₂₋₆alkenyl substituted with one or more substituents
each independently selected from halo, cyano or -C(=O)-NR⁹R¹⁰; or C₂₋₆alkynyl
substituted with one or more substituents each independently selected from halo,
15 cyano, -C(=O)-NR⁹R¹⁰;
- (i') X₃ is -NR⁵-or-O-;
- (j') R⁴ is halo, hydroxy, C₁₋₆alkyl, C₂₋₆alkenyl, C₂₋₆alkynyl, C₁₋₆alkyloxy, cyano, nitro,
-C(=O)-NR¹³R¹⁴, C₁₋₆alkyloxycarbonyl, C₁₋₆alkylcarbonyl, formyl, -NR¹³R¹⁴;
- (k') R⁵ is hydrogen; C₁₋₆alkyl;
- 20 (l') R⁶ is C₁₋₄alkyl;
- (m') R⁷ is any of the specific monocyclic or bicyclic, partially saturated or aromatic
carbocycles or monocyclic or bicyclic, partially saturated or aromatic heterocycles
specifically mentioned in this specification, wherein each of said carbocyclic or
heterocyclic ring systems may optionally be substituted with one, two or three
25 substituents each independently selected from halo, hydroxy, mercapto, C₁₋₆alkyl,
hydroxyC₁₋₆alkyl, aminoC₁₋₆alkyl, C₁₋₆alkylcarbonyl, C₁₋₆alkyloxy, C₁₋₆alkyloxy-
carbonyl, C₁₋₆alkylthio, cyano, nitro, polyhaloC₁₋₆alkyl, polyhaloC₁₋₆alkyloxy or
aminocarbonyl;
- (n') R⁸ is hydrogen or C₁₋₄alkyl;
- 30 (o') R⁹ and R¹⁰ each independently are hydrogen or C₁₋₆alkyl;
- (p') R¹³ and R¹⁴ each independently are hydrogen or C₁₋₆alkyl;
- (q') R¹⁵ is C₁₋₆alkyl optionally substituted with cyano or -C(=O)-NR¹³R¹⁴;
- (r') R¹⁷ is cyano, halo, hydroxy, -C(=O)-NR¹³R¹⁴, C₁₋₆alkyl optionally substituted with
cyano, -C(=O)-NR¹³R¹⁴; C₂₋₆alkenyl optionally substituted with cyano or
35 -C(=O)-NR¹³R¹⁴; C₂₋₆alkynyl optionally substituted with cyano or
-C(=O)-NR¹³R¹⁴; and, where possible, R¹⁷ may also be attached to the -b¹-b²-b³-

-15-

moiety by a double bond whereby R^{17} is then $=O$, $=NH$, $=N-R^{15}$, $=N-R^7$, $=N-O-R^{15}$, $=N-O-R^7$, $=CH_2$, $=CH-C(=O)-NR^{13}R^{14}$, $=CH-R^7$, or $=CH-R^{15}$; wherein $=CH_2$ may optionally be substituted with cyano;

(s') Q represents hydrogen or C_{1-6} alkyl or $-NR^9R^{10}$;

5 (t') Y represents hydrogen, hydroxy, halo, C_{1-6} alkyl, C_{1-6} alkyloxy, cyano, nitro, $NR^{13}R^{14}$, polyhalomethyloxy, $-NH-SO_2-R^8$, $-NH-SO_2-(C_{1-4}alkanediyl)-CO-N(R^8)_2$;

(u') aryl is phenyl or phenyl substituted with one, two or three substituents each independently selected from halo, hydroxy, C_{1-6} alkyl, hydroxy C_{1-6} alkyl, C_{1-6} alkylcarbonyl, C_{1-6} alkyloxy, C_{1-6} alkyloxycarbonyl, C_{1-6} alkylthio, cyano, nitro,
10 $-C(=O)-NR^{13}R^{14}$;

(v') Het is a monocyclic or bicyclic, partially saturated or aromatic heterocycle, specifically mentioned in this specification, wherein each of said heterocyclic ring systems may optionally be substituted with one, two or three substituents each independently selected from halo, hydroxy, mercapto, C_{1-6} alkyl, hydroxy C_{1-6} alkyl,
15 amino C_{1-6} alkyl, C_{1-6} alkylcarbonyl, C_{1-6} alkyloxy, C_{1-6} alkyloxycarbonyl, C_{1-6} alkylthio, cyano, nitro, polyhalo C_{1-6} alkyl, polyhalo C_{1-6} alkyloxy.

A specific subgroup of the compounds of formula (I) are those wherein all of the limitations (a') - (v') of the previous paragraph apply.

20

Of particular interest are any subgroups of the compounds of formula (I) wherein one or more of the afore mentioned limitations (a) - (v) or of the limitations (a') - (v') optionally apply as well as one or more of the following limitations (a'') - (v''):

25 (a'') $-a^1=a^2-a^3=a^4-$ represents a bivalent radical of formula



(b'') n is 1;

(c'') m is 1;

(d'') R^1 is hydrogen; methyl;

30 (e'') R^2 is halo, C_{1-6} alkyl optionally substituted with cyano, C_{2-6} alkenyl optionally substituted with cyano, C_{2-6} alkynyl optionally substituted with cyano, C_{1-6} alkyloxycarbonyl, carboxyl, cyano, amino, mono(C_{1-6} alkyl)amino, di(C_{1-6} alkyl)amino;

(f'') X_1 is $-NR^5-$, $-O-$, $-NR^{13}-C(=O)-$, $-C(=O)-NR^{13}-$;

35 (h'') R^3 is hydrogen, halo, C_{1-6} alkyl, $NR^{13}R^{14}$, $-C(=O)-NR^{13}R^{14}$, $-C(=O)-R^{15}$; C_{1-6} alkyl substituted with cyano; C_{2-6} alkenyl substituted with cyano; or C_{2-6} alkynyl substituted with cyano;


-16-

- (j'') R⁴ is halo, hydroxy, C₁₋₆alkyl, C₂₋₆alkenyl, C₂₋₆alkynyl, C₁₋₆alkyloxy, cyano, nitro, -C(=O)-NR¹³R¹⁴, -NR¹³R¹⁴;
- (k'') R⁵ is hydrogen; C₁₋₆alkyl;
- (m'') R⁷ is any of the specific monocyclic or bicyclic, partially saturated or aromatic carbocycles or monocyclic or bicyclic, partially saturated or aromatic heterocycles specifically mentioned in this specification, wherein each of said carbocyclic or heterocyclic ring systems may optionally be substituted with one, two or three substituents each independently selected from halo, hydroxy, mercapto, C₁₋₆alkyl, hydroxyC₁₋₆alkyl, aminoC₁₋₆alkyl, C₁₋₆alkylcarbonyl, C₁₋₆alkyloxy, C₁₋₆alkyloxy-carbonyl, C₁₋₆alkylthio, cyano, nitro, polyhaloC₁₋₆alkyl, polyhaloC₁₋₆alkyloxy or aminocarbonyl;
- (n'') R⁸ is hydrogen or C₁₋₄alkyl;
- (o'') R⁹ and R¹⁰ are hydrogen;
- (p'') R¹³ and R¹⁴ are hydrogen;
- (q'') R¹⁵ is C₁₋₆alkyl optionally substituted with cyano;
- (r'') R¹⁷ is cyano, -C(=O)-NR¹³R¹⁴, C₁₋₆alkyl optionally substituted with cyano, -C(=O)-NR¹³R¹⁴; C₂₋₆alkenyl optionally substituted with cyano or -C(=O)-NR¹³R¹⁴; C₂₋₆alkynyl optionally substituted with cyano or -C(=O)-NR¹³R¹⁴; and, where possible, R¹⁷ may also be attached to the -b¹-b²-b³- moiety by a double bond whereby R¹⁷ is then =O, =NH, =N-R¹⁵, =N-R⁷, =N-O-R¹⁵, =N-O-R⁷, =CH₂, =CH-C(=O)-NR¹³R¹⁴, =CH-R⁷, or =CH-R¹⁵; wherein =CH₂ may optionally be substituted with cyano;
- (s'') Q represents hydrogen or -NR⁹R¹⁰;
- (t'') Y represents hydrogen, hydroxy, halo, C₁₋₆alkyl, C₁₋₆alkyloxy, cyano, NR¹³R¹⁴, -NH-SO₂-R⁸, -NH-SO₂-(C₁₋₄alkanediyl)-CO-N(R⁸)₂;
- (u'') aryl is phenyl or phenyl substituted with one, two or three substituents each independently selected from halo, hydroxy, C₁₋₆alkyl, C₁₋₆alkyloxy, C₁₋₆alkylthio, cyano, nitro;
- (v'') Het is a monocyclic or bicyclic, partially saturated or aromatic heterocycle, specifically mentioned in this specification, wherein each of said heterocyclic ring systems may optionally be substituted with one, two or three substituents each independently selected from halo, hydroxy, mercapto, C₁₋₆alkyl, hydroxyC₁₋₆alkyl, aminoC₁₋₆alkyl, C₁₋₆alkylcarbonyl, C₁₋₆alkyloxy, C₁₋₆alkyloxy-carbonyl, C₁₋₆alkylthio, cyano, nitro, polyhaloC₁₋₆alkyl, polyhaloC₁₋₆alkyloxy.
- A specific subgroup of the compounds of formula (I) are those wherein all of the limitations (a'') - (v'') of the previous paragraph apply.

-17-

Still further particular groups of compounds are those compounds of formula (I) wherein one or wherever possible more of the following conditions apply :

- (a-1) m is 0, 1 or 2, in particular 1 or 2, more in particular 2; and wherein the R⁴ substituents are placed in the ortho position in respect of the X₁ moiety;
- 5 (a-2) X₁ is linked to one of the carbon atoms in meta position of the carbonatoms common to both rings of the bicyclic ring system to which X₁ is connected.
- (a-3) where applicable n is 0; or where applicable n is 1 and the R² substituent is placed in position 4 (para position) in respect of the NR¹-linker;
- (a-4) R² is hydroxy, halo, C₁₋₆alkyl optionally substituted with cyano or with -C(=O)R⁶,
- 10 C₃₋₇cycloalkyl, C₂₋₆alkenyl optionally substituted with one or more halogen atoms or cyano, C₂₋₆alkynyl optionally substituted with one or more halogen atoms or cyano, C₁₋₆alkyloxycarbonyl, carboxyl, cyano, nitro, amino, mono- or di(C₁₋₆alkyl)amino, polyhalomethyl, polyhalomethylthio, -S(=O)_pR⁶, -NH-S(=O)_pR⁶, -NHC(=O)H, -C(=O)NHNH₂, -NHC(=O)R⁶, -C(=NH)R⁶ or a radical of formula



15

wherein each A₁ independently is N, CH or CR⁶; and
A₂ is NH, O, S or NR⁶.

20

(a-5) R³ is R³ is hydrogen or C₁₋₆alkyl or C₁₋₆alkyl optionally substituted with cyano.

A preferred subgroup is that wherein R² is cyano and R¹ is hydrogen.

Also an interesting group of compounds are those compounds of formula (I) wherein one or more, preferably all of the following restrictions apply:

- 25 (b-1) n is at least 1, in particular 1; or n is 0;
- (b-2) R² is cyano;
- (b-3) m is 1, 2 or 3;
- (b-4) R⁴ is C₁₋₆alkyl, especially methyl; halo;
- (b-5) X₁ is NH or O;
- 30 (b-6) R¹ is hydrogen or C₁₋₄alkyl.

Of specific interest are those compounds of formula (I) or any of the subgroups specified herein, wherein R⁴ is halogen.

- 35 Also of specific interest are those compounds of formula (I) or any of the subgroups specified herein, wherein R¹⁷ is halo, cyano.

-18-

Another interesting group of compounds are those compounds of formula (I) or any of the subgroups specified herein, wherein R^{17} is oxo, C_{1-6} alkyl optionally substituted with cyano, $=N-O-C_{1-6}$ alkyl-Aryl, hydrogen, oxo, C_{1-6} alkyl optionally substituted with cyano or Het.

5

Further subgroups of the compounds in accordance with the present invention are those compounds of formula (I) or any of the subgroups of compounds of formula (I) specified herein, wherein one or more of C_{1-6} alkyl is limited to C_{1-4} alkyl, one or more of C_{1-4} alkyl is limited to C_{1-2} alkyl; wherein one or more of C_{2-6} alkenyl is limited to C_{2-4} alkenyl; wherein one or more of C_{2-6} alkynyl is limited to C_{2-4} alkynyl.

10

Further subgroups of the compounds in accordance with the present invention are those compounds of formula (I) or any of the subgroups of compounds of formula (I) specified herein, wherein one or more of the radicals that are (or one or more of the radicals that contain) heterocycles or carbocycles are the heterocycles or carbocycles as specifically set forth therein.

15


Synthesis

The compounds of formula (I) can be prepared via a number of pathways a number of which are explained hereinafter in more detail.

20

The compounds of formula (I) can be generally prepared by reacting an indane (a-1) with pyrimidine derivative (a-2). The groups HX_1 and W_1 are selected such that a X_1 linking moiety is formed.

25



-19-

In particular, W_1 is a suitable leaving group and X_1 is a heteroatom. Examples of suitable leaving groups in (a-2) are halogen, in particular chloro and bromo, tosylate, mesylate, triflate and the like groups. For the preparation of compounds of formula (I) wherein X_1 is other than a heteroatom, $-W_1$ and $-X_1H$ can have other meanings as outlined hereinafter.

The conversion of (a-1) with (a-2) to (a-3) in the above scheme is particularly useful when W_1 is a leaving group and X_1 is a heteroatom such as $-NR^5$ -, $-NH-NH$ -, $-N=N$ -, $-O$ -, $-S$ -, $-X_2-C_{1-4}alkanediy-$. This conversion is particularly suited in the instance where X_1 is $-O$ -. In the instance where is S , the latter can conveniently be transferred to the corresponding sulfoxide or sulfon using art-known oxidation procedures.

The above reaction usually is performed in the presence of a suitable solvent. Suitable solvents are for example acetonitrile, alcohols, such as for example ethanol, 2-propanol, ethylene glycol, propylene glycol, polar aprotic solvents such as *N,N*-dimethylformamide; *N,N*-dimethylacetamide, dimethylsulfoxide, 1-methyl-2-pyrrolidinone, [bmim]PF₃; ethers such as 1,4-dioxane, propylene glycol monomethylether.

Where X_1 is $-C(=O)-$ a starting material (a-1) wherein the group $-X_1H$ is a Grignard type of group ($-Mg-halo$) or lithium is reacted with a starting material (a-2) wherein W_1 is an ester ($-COOalkyl$). The latter ester may also be reduced to an alcohol with e.g. $LiAlH_4$ and subsequently oxidized with a mild oxidant such as MnO_2 to the corresponding aldehyde which subsequently is reacted with the (a-1) starting material wherein the group $-X_1H$ is a Grignard type of group ($-Mg-halo$) or lithium. The compounds wherein $-X_1-$ is $-C(=O)-$ can be converted to the $-CHOH-$ analogs by a suitable reduction reaction e.g. with $LiAlH_4$.

Where X_1 is $C_{1-4}alkanediy-$ the linkage can be introduced by a Grignard reaction, e.g. by reacting a starting material (a-1) wherein the $-X_1H$ group is $-C_{1-4}alkanediy-Mg-halo$ with an (a-2) reagent wherein W_1 is a halo group, or vice versa. Where X_1 is methylene, the methylene group can be oxidized to a $-C(=O)-$ group (X_1 is $-C(=O)-$) e.g. with selenium dioxide. The $-C(=O)-$ group in turn can be reduced with a suitable hydride such as $LiAlH_4$ to a $-CHOH-$ group.

Where X_1 is $-NR^{13}-C(=O)-$, or $-C(=O)-NR^{13}-$, the X_1 linkage can be formed via a suitable amide bond forming reaction starting from an intermediate (a-1) wherein $-X_1H$ is $-NHR^{13}$ and an intermediate (a-2) wherein W_1 is a carboxyl group or an active derivative thereof, or vice versa starting from an intermediate (a-1) wherein $-X_1H$ is carboxyl group or an active derivative thereof and an intermediate (a-2) wherein W_1 is a

-20-

group $-NHR^{13}$. The amide bond may be formed following methodologies generally known in the art, e.g. by activation of the carboxyl group to a carbonyl chloride or bromide or by using a suitable coupling agent.

- 5 Where X_1 is $-X_2-C_{1-4}alkanedyl-$, an intermediate (a-1) wherein the group $-X_1H$ is a radical $-X_2H$ is reacted with an intermediate (a-2) wherein the group W_1 is $-C_{1-4}alkanedyl-W_2$, wherein W_2 in turn is a suitable leaving group. Or where X_1 is $-C_{1-4}alkanedyl-X_2-$ an intermediate (a-1) wherein the group $-X_1H$ is $-C_{1-4}alkanedyl-W_2$, wherein W_2 in turn is a suitable leaving group, is reacted with an intermediate (a-2) wherein W_1 is $-X_2H$.

The linkages of X_2 being other than a heteroatom (i.e. X_2 is $-C(=O)-$, $-CHOH-$) can be prepared by analogous procedures as for introducing the linker X_1 .

- 15 In the instance where X_1 is $-NR^5-$ the reaction of (a-1) with reagent (a-2) is typically conducted under neutral conditions or, which is preferred, under acidic conditions, usually at elevated temperatures and under stirring. The acid conditions may be obtained by adding amounts of a suitable acid or by using acidic solvents, e.g. hydrochloric acid dissolved in an alkanol such as 1- or 2-propanol or in acetonitrile.

20 The above reaction can be performed in the presence of a suitable solvent. Suitable solvents are for example acetonitrile, an alcohol, such as for example ethanol, 2-propanol, 2-propanol-HCl; *N,N*-dimethylformamide; *N,N*-dimethylacetamide, 1-methyl-2-pyrrolidinone; 1,4-dioxane, propyleneglycol monomethylether.

- 25 Preferably the solvent is 2-propanol, 6 N HCl in 2-propanol or acetonitrile, especially acetonitrile. Optionally, sodium hydride may be present.

In the instance where X_1 is $-O-$, the reaction is typically conducted as follows.

- 30 Intermediate (a-1) is first reacted under stirring at room temperature with hydrides in an organic solvent. Subsequently, a solvent, such as *N*-methylpyrrolidinone, dimethylacetamide or dimethylformamide, is added to the mixture and followed by the addition of reagent (a-2). Typically, the reaction mixture is stirred overnight at elevated temperatures to yield compound (a-3).


- 35 The compounds of formula (a-3) having a R^{17} substituent which is an oxo ($=O$) group (represented by structure (a-3-1)) can be used as a starting material to obtain compounds of formula (I) having a R^{17} substituent which is a $=N-R^{18}$ substituent, wherein $=N-R^{18}$ is $=NH$, $=N-R^{15}$, $=N-R^7$, $=N-O-R^{15}$, $=N-O-R^7$ as defined above. In this reaction pathway, intermediate (a-3-1) is reacted with reagent (a-7) (reagent (a-7) is of

-21-

general formula NH_3 , $\text{NH}_2\text{-R}^{15}$, $\text{NH}_2\text{-R}^7$, $\text{NH}_2\text{-O-R}^{15}$, $\text{NH}_2\text{-O-R}^7$, in particular Aryl- $\text{C}_{1-6}\text{alkyl-O-NH}_2$) at elevated temperatures in an alcoholic solvent in the presence of a base to generate a compound of formula (a-8).

- 5 Similarly, the compounds of formula (a-3-1) can be used as a starting material to obtain compounds of formula (I) having a R^{17} substituent which is a $=\text{X}$ substituent, wherein $=\text{X}$ is $=\text{CH}_2$, $=\text{CH-C(=O)-NR}^{13}\text{R}^{14}$, $=\text{CH-R}^7$, or $=\text{CH-R}^{15}$ as defined above. Intermediate (a-3) is further reacted with reagent (a-4) in a Wittig reaction or a Wittig-Horner reaction. In the former instance, reagent (a-4) is a Wittig type reagent, such as a
- 10 triphenylphosphoniumylide, in the latter instance, a Wittig-Horner type of reagent, in particular a phosphonate, such as e.g. a reagent of formula $\text{di(C}_{1-6}\text{alkyloxy)-P(=O)-X}_4$, wherein X_4 is a substituent R^{17} that can be linked to the ring via a double bond (exo double bond). The Wittig-Horner type of conversion typically is conducted in the presence of a base, preferably a strong base, in an aprotic organic solvent at room
- 15 temperature. The reaction should be allowed sufficient time to complete, typically it is allowed to proceed overnight to yield compound (a-5). This latter compound may further be reacted in an alcoholic solvent under reducing conditions to generate a compound of formula (a-6).
- 20 Both conversion reactions are outlined in the following reaction scheme.

-22-




The oxo group in the compounds (a-3-1) may also be at other positions in the ring having the R^{17} substituent(s), the same type of derivatisation may be performed resulting in the topical isomers of (a-8), (a-5) and (a-6).

5

The compounds of formula (I) can also be prepared as outlined in the reaction scheme here below.

-23-



An indane of formula (a-1) is reacted with a pyrimidine (a-9) wherein the substituents have the meanings specified herein and W_1 is a suitable leaving group such as, for example, halo, triflate, tosylate, methylsulfonyl and the like, yielding an intermediate (a-10). This reaction can be done similarly as outlined above for the reaction of (a-1) with (a-2), in particular for the various possibilities of the linker $-X_1-$. Where necessary, the W_1 group that does not intervene in this reaction may be replaced by a leaving group precursor such as a OH functionality which a particular stage of the reaction procedure is converted to a leaving group, e.g. by converting the OH group into a halogen group, or by reacting it with a suitable reagent such as $POCl_3$, tosyl chloride, mesyl chloride and the like

The end products (I) can be prepared from this starting material (a-10) by reaction with the amino substituted aromatic compound (a-11) in an arylation type of reaction.

Suitable solvents for the reaction of (a-1) with (a-9) and of (a-10) with (a-11) are ethers, e.g. 1,4-dioxane, THF, alcohols, ethanol, propanol, butanol, ethylene glycol, propylene glycol, propylene glycol monomethyl ether, the aprotic solvents such as acetonitrile, DMF, DMA, 1-methyl-2-pyrrolidinone and the like. If necessary a base can be added. Suitable bases in this reaction are for example sodium acetate, potassium acetate, *N,N*-diethylethanamine, sodium hydrogencarbonate, sodium hydroxide and the like.

In this and the following preparations, the reaction products may be isolated from the reaction medium and, if necessary, further purified according to methodologies generally known in the art such as, for example, extraction, crystallization, distillation, trituration and chromatography.

5

The compounds of formula (I) may further be prepared by converting compounds of formula (I) into each other according to art-known group transformation reactions.

10

The compounds of formula (I) may be converted to the corresponding *N*-oxide forms following art-known procedures for converting a trivalent nitrogen into its *N*-oxide form. Said *N*-oxidation reaction may generally be carried out by reacting the starting material of formula (I) with an appropriate organic or inorganic peroxide. Appropriate inorganic peroxides comprise, for example, hydrogen peroxide, alkali metal or earth alkaline metal peroxides, e.g. sodium peroxide, potassium peroxide; appropriate organic peroxides may comprise peroxy acids such as, for example, benzenecarboperoxoic acid or halo substituted benzenecarboperoxoic acid, e.g. 3-chlorobenzenecarboperoxoic acid, peroxyalkanoic acids, e.g. peroxyacetic acid, alkylhydroperoxides, e.g. tert.butyl hydroperoxide. Suitable solvents are, for example, water, lower alcohols, e.g. ethanol and the like, hydrocarbons, e.g. toluene, ketones, e.g. 2-butanone, halogenated hydrocarbons, e.g. dichloromethane, and mixtures of such solvents.

15

20

For instance, a compound of formula (I) wherein R^3 comprises cyano, can be converted into a compound of formula (I) wherein R^3 comprises aminocarbonyl, by reaction with $HCOOH$, in the presence of a suitable acid, such as hydrochloric acid. A compound of formula (I) wherein R^3 comprises cyano, can also further be converted into a compound of formula (I) wherein R^3 comprises tetrazolyl, by reaction with sodium azide in the presence of ammonium chloride and *N,N*-dimethylacetamide.

25

Compounds of formula (I) wherein R^3 comprises aminocarbonyl, can be converted into a compound of formula (I) wherein R^3 comprises cyano, in the presence of a suitable dehydrating agent. The dehydration can be performed according to methodologies well-known to the person skilled in the art, such as the ones disclosed in "Comprehensive Organic Transformations. A guide to functional group preparations" by Richard C. Larock, John Wiley & Sons, Inc, 1999, p 1983-1985, which is incorporated herein as reference. Different suitable reagents are enumerated in said reference, such as for example $SOCl_2$, $HOSO_2NH_2$, $ClSO_2NCO$, $MeO_2CNSO_2NEt_3$, $PhSO_2Cl$, $TsCl$, P_2O_5 , $(Ph_3PO_3SCF_3)_3O_3SCF_3$, polyphosphate ester, $(EtO)_2POP(OEt)_2$, $(EtO)_3PI_2$, 2-chloro-1,3,2-dioxaphospholane, 2,2,2-trichloro-2,2-dihydro-1,3,2-dioxaphospholane, $POCl_3$, PPh_3 , $P(NCl_2)_3$, $P(NEt_2)_3$, $COCl_2$, $NaCl$, $AlCl_3$, $ClCOCOC$,

30

35

-25-

ClCO_2Me , Cl_3CCOCl , $(\text{CF}_3\text{CO})_2\text{O}$, $\text{Cl}_3\text{CN}=\text{CCl}_2$, 2,4,6-trichloro-1,3,5-triazine, $\text{NaCl}\cdot\text{AlCl}_3$, $\text{HN}(\text{SiMe}_2)_3$, $\text{N}(\text{SiMe}_2)_4$, LiAlH_4 and the like. All the reagents listed in said publication are incorporated herein as reference.

- 5 Compounds of formula (I) wherein R^3 comprises C_{2-6} alkenyl can be converted into a compound of formula (I) wherein R^3 comprises C_{1-6} alkyl by reduction in the presence of a suitable reducing agent, such as for example H_2 , in the presence of a suitable catalyst, such as for example palladium on charcoal, and in the presence of a suitable solvent, such as for example an alcohol, e.g. methanol.

10

Compounds of formula (I) wherein R^3 represents $\text{CH}(\text{OH})\text{-R}^{16}$, can be converted into a compound of formula (I) wherein R^3 represents $\text{C}(=\text{O})\text{-R}^{16}$ by reaction with Jones's reagent in the presence of a suitable solvent, such as for example 2-propanone.

- 15 Compound of formula (I) wherein R^3 represents $\text{C}(=\text{O})\text{-CH}_2\text{-R}^{16a}$, wherein R^{16a} represents cyano or aminocarbonyl, can be converted into a compound of formula (I) wherein R^3 represents $\text{C}(\text{Cl})=\text{CH-R}^{16a}$ by reaction with POCl_3 .

- 20 Compounds of formula (I) wherein R^3 represents a monocyclic, bicyclic or tricyclic saturated, partially saturated or aromatic carbocycle or a monocyclic, bicyclic or tricyclic saturated, partially saturated or aromatic heterocycle substituted with formyl can be converted into compounds of formula (I) wherein R^3 represents a monocyclic, bicyclic or tricyclic saturated, partially saturated or aromatic carbocycle or a monocyclic, bicyclic or tricyclic saturated, partially saturated or aromatic heterocycle substituted with $\text{CH}(=\text{N-O-R}^8)$ by reaction with NH_2OR^8 in the presence of a suitable base, such as for example sodium hydroxide and a suitable solvent, such as for example an alcohol, e.g. ethanol and the like. Compounds of formula (I) wherein R^3 represents a monocyclic, bicyclic or tricyclic saturated, partially saturated or aromatic carbocycle or a monocyclic, bicyclic or tricyclic saturated, partially saturated or aromatic heterocycle substituted with $\text{CH}(=\text{N-O-R}^8)$ can be converted into a compound of formula (I) wherein R^3 represents a monocyclic, bicyclic or tricyclic saturated, partially saturated or aromatic carbocycle or a monocyclic, bicyclic or tricyclic saturated, partially saturated or aromatic heterocycle substituted with CN by reaction with a carbodiimide in the presence of a suitable solvent, such as for example tetrahydrofuran.

35

Compounds of formula (I) wherein R^4 represents nitro, can be converted into a compound of formula (I) wherein R^4 is amino, in the presence of a suitable reducing agent, such as for example H_2 , in the presence of a suitable catalyst, such as for example

Raney Nickel, and in the presence of a suitable solvent, such as for example an alcohol, e.g. methanol.

5 Compounds of formula (I) wherein R^1 is hydrogen, can be converted into a compound of formula (I) wherein R^1 is C_{1-6} alkyl, by reaction with a suitable alkylating agent, such as for example iodo- C_{1-6} alkyl, in the presence of a suitable base, such as for example sodium hydride, and a suitable solvent, such as for example tetrahydrofuran.

10 Compounds of formula (I) having a carbon-carbon double bond can be reduced to the corresponding compounds with a single bond using catalytic hydrogenation procedures. In these procedures use is made of a noble metal catalyst. An attractive such catalyst is Pd. The palladium (Pd) catalyst may be a homogeneous Pd catalyst, such as for example $Pd(OAc)_2$, $PdCl_2$, $Pd(PPh_3)_4$, $Pd(PPh_3)_2Cl_2$, bis(dibenzylidene acetone) palladium, palladium thiomethylphenylglutaramide metallacycle and the like, or a heterogeneous Pd catalyst, such as for example palladium on charcoal, palladium on metal oxides, 15 palladium on zeolites.

20 Preferably, the palladium catalyst is a heterogeneous Pd catalyst, more preferably palladium on charcoal (Pd/C). Pd/C is a recoverable catalyst, is stable and relatively inexpensive. It can be easily separated (filtration) from the reaction mixture thereby reducing the risk of Pd traces in the final product. The use of Pd/C also avoids the need for ligands, such as for example phosphine ligands, which are expensive, toxic and contaminants of the synthesized products.


25 Some of the compounds of formula (I) and some of the intermediates in the present invention may contain an asymmetric carbon atom. Pure stereochemically isomeric forms of said compounds and said intermediates can be obtained by the application of art-known procedures. For example, diastereoisomers can be separated by physical methods such as selective crystallization or chromatographic techniques, e.g. counter 30 current distribution, liquid chromatography and the like methods. Enantiomers can be obtained from racemic mixtures by first converting said racemic mixtures with suitable resolving agents such as, for example, chiral acids, to mixtures of diastereomeric salts or compounds; then physically separating said mixtures of diastereomeric salts or compounds by, for example, selective crystallization or chromatographic techniques, 35 e.g. liquid chromatography and the like methods; and finally converting said separated diastereomeric salts or compounds into the corresponding enantiomers. Pure stereochemically isomeric forms may also be obtained from the pure stereochemically isomeric forms of the appropriate intermediates and starting materials, provided that the intervening reactions occur stereospecifically.

-27-

An alternative manner of separating the enantiomeric forms of the compounds of formula (I) and intermediates involves liquid chromatography, in particular liquid chromatography using a chiral stationary phase.

Some of the intermediates and starting materials are known compounds and may be commercially available or may be prepared according to art-known procedures or some of the compounds of formula (I) or the described intermediates may be prepared according to the procedures described in WO 99/50250 and WO 00/27825.

Intermediates of formula (a-2) wherein W_1 represents a leaving group, can be prepared by reacting an intermediate of formula (b-1) with a suitable halogenating agent, e.g. N-bromosuccinimide, N-chlorosuccinimide, PCl_3 , PCl_5 or with a suitable leaving group introducing agent of formula (b-2) wherein W_1 represents the leaving group and L represents part of the leaving group introducing agent, such as for example $POCl_3$, triflyl chloride, tosyl chloride, mesyl chloride and the like.



This reaction typically is conducted in a suitable solvent, if desired, in the presence of a suitable base, for example, sodium acetate, potassium acetate, N,N -diethylethanamine, sodium hydrogencarbonate, sodium hydroxide and the like.

Suitable solvents in the above reaction are for example acetonitrile, N,N -dimethylacetamide, an ionic liquid e.g. [bmim] PF_6 , N,N -dimethylformamide, water, tetrahydrofuran, dimethylsulphoxide, 1-methyl-2-pyrrolidinone and the like.

The compounds of formula (I) as prepared in the hereinabove described processes may be synthesized as a mixture of stereoisomeric forms, in particular in the form of racemic mixtures of enantiomers which can be separated from one another following art-known resolution procedures. The racemic compounds of formula (I) may be converted into the corresponding diastereomeric salt forms by reaction with a suitable chiral acid. Said diastereomeric salt forms are subsequently separated, for example, by selective or fractional crystallization and the enantiomers are liberated therefrom by alkali. An alternative manner of separating the enantiomeric forms of the compounds of formula (I) involves liquid chromatography using a chiral stationary phase. Said pure

-28-

stereochemically isomeric forms may also be derived from the corresponding pure stereochemically isomeric forms of the appropriate starting materials, provided that the reaction occurs stereospecifically. Preferably if a specific stereoisomer is desired, said compound will be synthesized by stereospecific methods of preparation. These methods
5 will advantageously employ enantiomerically pure starting materials.

It will be appreciated by those skilled in the art that in the processes described above the functional groups of intermediate compounds may need to be blocked by protecting
10 groups.

Functional groups, which are desirable to protect include hydroxy, amino and carboxylic acid. Suitable protecting groups for hydroxy include trialkylsilyl groups (e.g. *tert*-butyldimethylsilyl, *tert*-butyldiphenylsilyl or trimethylsilyl), benzyl and tetrahydropyran-
15 benzyloxycarbonyl. Suitable protecting groups for amino include *tert*-butyloxycarbonyl or benzyloxycarbonyl. Suitable protecting groups for carboxylic acid include C₁₋₆alkyl or benzyl esters.

The protection and deprotection of functional groups may take place before or after a
20 reaction step.

The use of protecting groups is fully described in 'Protective Groups in Organic Chemistry', edited by J W F McOmie, Plenum Press (1973), and 'Protective Groups in Organic Synthesis' 2nd edition, T W Greene & P G M Wutz, Wiley Interscience (1991).

25 The compounds of formula (I) show antiretroviral properties (reverse transcriptase inhibiting properties), in particular against Human Immunodeficiency Virus (HIV), which is the aetiological agent of Acquired Immune Deficiency Syndrome (AIDS) in humans. The HIV virus preferentially infects human T-4 cells and destroys them or changes their normal function, particularly the coordination of the immune system. As a
30 result, an infected patient has an ever decreasing number of T-4 cells, which moreover behave abnormally. Hence, the immunological defense system is unable to combat infections and neoplasms and the HIV infected subject usually dies by opportunistic infections such as pneumonia, or by cancers. Other conditions associated with HIV infection include thrombocytopaenia, Kaposi's sarcoma and infection of the central
35 nervous system characterized by progressive demyelination, resulting in dementia and symptoms such as, progressive dysarthria, ataxia and disorientation. HIV infection further has also been associated with peripheral neuropathy, progressive generalized lymphadenopathy (PGL) and AIDS-related complex (ARC).

The present compounds also show activity against (multi) drug resistant HIV strains, in particular (multi) drug resistant HIV-1 strains, more in particular the present compounds show activity against HIV strains, especially HIV-1 strains, that have acquired resistance to one or more art-known non-nucleoside reverse transcriptase inhibitors. Art-known
5 non-nucleoside reverse transcriptase inhibitors are those non-nucleoside reverse transcriptase inhibitors other than the present compounds and in particular commercial non-nucleoside reverse transcriptase inhibitors. The present compounds also have little or no binding affinity to human α -1 acid glycoprotein; human α -1 acid glycoprotein does not or only weakly affect the anti HIV activity of the present compounds.

10 Due to their antiretroviral properties, particularly their anti-HIV properties, especially their anti-HIV-1-activity, the compounds of formula (I), their *N*-oxides, pharmaceutically acceptable addition salts, quaternary amines and stereochemically isomeric forms thereof, are useful in the treatment of individuals infected by HIV and for
15 the prophylaxis of these infections. In general, the compounds of the present invention may be useful in the treatment of warm-blooded animals infected with viruses whose existence is mediated by, or depends upon, the enzyme reverse transcriptase. Conditions which may be prevented or treated with the compounds of the present invention, especially conditions associated with HIV and other pathogenic retroviruses, include
20 AIDS, AIDS-related complex (ARC), progressive generalized lymphadenopathy (PGL), as well as chronic Central Nervous System diseases caused by retroviruses, such as, for example HIV mediated dementia and multiple sclerosis.

Thus in a further aspect, the compounds of the present invention including any subgroup
25 defined herein may therefore be used as a medicine in particular against above-mentioned conditions. Said use as a medicine or method of treatment comprises the administration to HIV-infected subjects of an amount effective to combat the conditions associated with HIV and other pathogenic retroviruses, especially HIV-1. In particular, the compounds of formula (I) may be used in the manufacture of a medicament for the
30 treatment or the prevention of HIV infections.

In view of the pharmacological properties of the compounds of formula (I), there is provided a method of treating warm-blooded animals, including humans, suffering from, or a method of preventing warm-blooded animals, including humans, to suffer from,
35 viral infections, especially HIV infections. Said method comprises the administration, preferably oral administration, of an effective amount of a compound of formula (I), a *N*-oxide form, a pharmaceutically acceptable addition salt, a quaternary amine or a possible stereoisomeric form thereof, to warm-blooded animals, including humans.

-30-

In another aspect, the present invention also provides pharmaceutical compositions comprising a therapeutically effective amount of a compound of formula (I) and a pharmaceutically acceptable carrier or diluent. In still a further aspect there is provided a method of preparation a pharmaceutical composition as specified herein comprising mixing a compound of formula (I) with a suitable pharmaceutically acceptable carrier or diluent.

The compounds of the present invention or any subgroup thereof may be formulated into various pharmaceutical forms for administration purposes. As appropriate compositions there may be cited all compositions usually employed for systemically administering drugs. To prepare the pharmaceutical compositions of this invention, an effective amount of the particular compound, optionally in addition salt form, as the active ingredient is combined in intimate admixture with a pharmaceutically acceptable carrier, which carrier may take a wide variety of forms depending on the form of preparation desired for administration. These pharmaceutical compositions are desirable in unitary dosage form suitable, particularly, for administration orally, rectally, percutaneously, or by parenteral injection. For example, in preparing the compositions in oral dosage form, any of the usual pharmaceutical media may be employed such as, for example, water, glycols, oils, alcohols and the like in the case of oral liquid preparations such as suspensions, syrups, elixirs, emulsions and solutions; or solid carriers such as starches, sugars, kaolin, diluents, lubricants, binders, disintegrating agents and the like in the case of powders, pills, capsules, and tablets. Because of their ease in administration, tablets and capsules represent the most advantageous oral dosage unit forms, in which case solid pharmaceutical carriers are obviously employed. For parenteral compositions, the carrier will usually comprise sterile water, at least in large part, though other ingredients, for example, to aid solubility, may be included. Injectable solutions, for example, may be prepared in which the carrier comprises saline solution, glucose solution or a mixture of saline and glucose solution. Injectable suspensions may also be prepared in which case appropriate liquid carriers, suspending agents and the like may be employed. Also included are solid form preparations which are intended to be converted, shortly before use, to liquid form preparations. In the compositions suitable for percutaneous administration, the carrier optionally comprises a penetration enhancing agent and/or a suitable wetting agent, optionally combined with suitable additives of any nature in minor proportions, which additives do not introduce a significant deleterious effect on the skin. Said additives may facilitate the administration to the skin and/or may be helpful for preparing the desired compositions. These compositions may be administered in various ways, e.g., as a transdermal patch, as a spot-on, as an ointment.

-31-

The compounds of the present invention may also be administered via inhalation or insufflation by means of methods and formulations employed in the art for administration via this way. Thus, in general the compounds of the present invention may be administered to the lungs in the form of a solution, a suspension or a dry powder. Any system developed for the delivery of solutions, suspensions or dry powders via oral or nasal inhalation or insufflation are suitable for the administration of the present compounds.

To aid solubility of the compounds of formula (I), suitable ingredients, e.g. cyclodextrins, may be included in the compositions. Appropriate cyclodextrins are α -, β -, γ -cyclodextrins or ethers and mixed ethers thereof wherein one or more of the hydroxy groups of the anhydroglucose units of the cyclodextrin are substituted with C₁₋₆alkyl, particularly methyl, ethyl or isopropyl, e.g. randomly methylated β -CD; hydroxyC₁₋₆alkyl, particularly hydroxyethyl, hydroxy-propyl or hydroxybutyl; carboxyC₁₋₆alkyl, particularly carboxymethyl or carboxy-ethyl; C₁₋₆alkylcarbonyl, particularly acetyl. Especially noteworthy as complexants and/or solubilizers are β -CD, randomly methylated β -CD, 2,6-dimethyl- β -CD, 2-hydroxyethyl- β -CD, 2-hydroxyethyl- β -CD, 2-hydroxypropyl- β -CD and (2-carboxymethoxy)propyl- β -CD, and in particular 2-hydroxypropyl- β -CD (2-HP- β -CD).

The term mixed ether denotes cyclodextrin derivatives wherein at least two cyclodextrin hydroxy groups are etherified with different groups such as, for example, hydroxy-propyl and hydroxyethyl.

The average molar substitution (M.S.) is used as a measure of the average number of moles of alkoxy units per mole of anhydroglucose. The average substitution degree (D.S.) refers to the average number of substituted hydroxyls per anhydroglucose unit. The M.S. and D.S. value can be determined by various analytical techniques such as nuclear magnetic resonance (NMR), mass spectrometry (MS) and infrared spectroscopy (IR). Depending on the technique used, slightly different values may be obtained for one given cyclodextrin derivative. Preferably, as measured by mass spectrometry, the M.S. ranges from 0.125 to 10 and the D.S. ranges from 0.125 to 3.

Other suitable compositions for oral or rectal administration comprise particles consisting of a solid dispersion comprising a compound of formula (I) and one or more appropriate pharmaceutically acceptable water-soluble polymers.

The term "a solid dispersion" used hereinafter defines a system in a solid state (as opposed to a liquid or gaseous state) comprising at least two components, in casu the

compound of formula (I) and the water-soluble polymer, wherein one component is dispersed more or less evenly throughout the other component or components (in case additional pharmaceutically acceptable formulating agents, generally known in the art, are included, such as plasticizers, preservatives and the like). When said dispersion of the components is such that the system is chemically and physically uniform or homogenous throughout or consists of one phase as defined in thermo-dynamics, such a solid dispersion will be called "a solid solution". Solid solutions are preferred physical systems because the components therein are usually readily bioavailable to the organisms to which they are administered. This advantage can probably be explained by the ease with which said solid solutions can form liquid solutions when contacted with a liquid medium such as the gastro-intestinal juices. The ease of dissolution may be attributed at least in part to the fact that the energy required for dissolution of the components from a solid solution is less than that required for the dissolution of components from a crystalline or microcrystalline solid phase.

The term "a solid dispersion" also comprises dispersions, which are less homogenous throughout than solid solutions. Such dispersions are not chemically and physically uniform throughout or comprise more than one phase. For example, the term "a solid dispersion" also relates to a system having domains or small regions wherein amorphous, microcrystalline or crystalline compound of formula (I), or amorphous, microcrystalline or crystalline water-soluble polymer, or both, are dispersed more or less evenly in another phase comprising water-soluble polymer, or compound of formula (I), or a solid solution comprising compound of formula (I) and water-soluble polymer. Said domains are regions within the solid dispersion distinctively marked by some physical feature, small in size, and evenly and randomly distributed throughout the solid dispersion.

Various techniques exist for preparing solid dispersions including melt-extrusion, spray-drying and solution-evaporation.

The solution-evaporation process comprises the following steps :

- a) dissolving the compound of formula (I) and the water-soluble polymer in an appropriate solvent, optionally at elevated temperatures;
- b) heating the solution resulting under point a), optionally under vacuum, until the solvent is evaporated. The solution may also be poured onto a large surface so as to form a thin film, and evaporating the solvent therefrom.

In the spray-drying technique, the two components are also dissolved in an appropriate solvent and the resulting solution is then sprayed through the nozzle of a spray dryer

-33-

followed by evaporating the solvent from the resulting droplets at elevated temperatures.

The preferred technique for preparing solid dispersions is the melt-extrusion process comprising the following steps :

- 5 a) mixing a compound of formula (I) and an appropriate water-soluble polymer,
- b) optionally blending additives with the thus obtained mixture,
- c) heating and compounding the thus obtained blend until one obtains a homogenous melt,
- 10 d) forcing the thus obtained melt through one or more nozzles; and
- e) cooling the melt until it solidifies.

20 The terms "melt" and "melting" should be interpreted broadly. These terms not only mean the alteration from a solid state to a liquid state, but can also refer to a transition to a glassy state or a rubbery state, and in which it is possible for one component of the mixture to get embedded more or less homogeneously into the other. In particular cases, one component will melt and the other component(s) will dissolve in the melt thus forming a solution, which upon cooling may form a solid solution having advantageous dissolution properties.

20 After preparing the solid dispersions as described hereinabove, the obtained products can be optionally milled and sieved.

25 The solid dispersion product may be milled or ground to particles having a particle size of less than 600 μm , preferably less than 400 μm and most preferably less than 125 μm .

 The particles prepared as described hereinabove can then be formulated by conventional techniques into pharmaceutical dosage forms such as tablets and capsules.

30 It will be appreciated that a person of skill in the art will be able to optimize the parameters of the solid dispersion preparation techniques described above, such as the most appropriate solvent, the working temperature, the kind of apparatus being used, the rate of spray-drying, the throughput rate in the melt-extruder

35 The water-soluble polymers in the particles are polymers that have an apparent viscosity, when dissolved at 20°C in an aqueous solution at 2 % (w/v), of 1 to 5000 mPa.s more preferably of 1 to 700 mPa.s, and most preferred of 1 to 100 mPa.s. For example, suitable water-soluble polymers include alkylcelluloses, hydroxyalkylcelluloses, hydroxyalkyl alkylcelluloses, carboxyalkylcelluloses, alkali metal salts of carboxyalkyl-

celluloses, carboxyalkylalkylcelluloses, carboxyalkylcellulose esters, starches, pectines, chitin derivates, di-, oligo- and polysaccharides such as trehalose, alginic acid or alkali metal and ammonium salts thereof, carrageenans, galactomannans, tragacanth, agar-agar, gummi arabicum, guar gummi and xanthan gummi, polyacrylic acids and the salts thereof, polymethacrylic acids and the salts thereof, methacrylate copolymers, polyvinylalcohol, polyvinylpyrrolidone, copolymers of polyvinylpyrrolidone with vinyl acetate, combinations of polyvinylalcohol and polyvinylpyrrolidone, polyalkylene oxides and copolymers of ethylene oxide and propylene oxide. Preferred water-soluble polymers are hydroxypropyl methylcelluloses.

10

Also one or more cyclodextrins can be used as water soluble polymer in the preparation of the above-mentioned particles as is disclosed in WO 97/18839. Said cyclodextrins include the pharmaceutically acceptable unsubstituted and substituted cyclodextrins known in the art, more particularly α , β or γ cyclodextrins or the pharmaceutically acceptable derivatives thereof.

15

Substituted cyclodextrins which can be used to prepare the above described particles include polyethers described in U.S. Patent 3,459,731. Further substituted cyclodextrins are ethers wherein the hydrogen of one or more cyclodextrin hydroxy groups is replaced by C₁-6alkyl, hydroxyC₁-6alkyl, carboxy-C₁-6alkyl or C₁-6alkyloxycarbonylC₁-6alkyl or mixed ethers thereof. In particular such substituted cyclodextrins are ethers wherein the hydrogen of one or more cyclodextrin hydroxy groups is replaced by C₁-3alkyl, hydroxyC₂-4alkyl or carboxyC₁-2alkyl or more in particular by methyl, ethyl, hydroxyethyl, hydroxypropyl, hydroxybutyl, carboxy-methyl or carboxyethyl.

20

25

Of particular utility are the β -cyclodextrin ethers, e.g. dimethyl- β -cyclodextrin as described in Drugs of the Future, Vol. 9, No. 8, p. 577-578 by M. Nogradi (1984) and polyethers, e.g. hydroxypropyl β -cyclodextrin and hydroxyethyl β -cyclodextrin, being examples. Such an alkyl ether may be a methyl ether with a degree of substitution of about 0.125 to 3, e.g. about 0.3 to 2. Such a hydroxypropyl cyclodextrin may for example be formed from the reaction between β -cyclodextrin and propylene oxide and may have a MS value of about 0.125 to 10, e.g. about 0.3 to 3.

30

Another type of substituted cyclodextrins is sulfobutylcyclodextrines.

35

The ratio of the compound of formula (I) over the water soluble polymer may vary widely. For example ratios of 1/100 to 100/1 may be applied. Interesting ratios of the compound of formula (I) over cyclodextrin range from about 1/10 to 10/1. More interesting ratios range from about 1/5 to 5/1.

-35-

It may further be convenient to formulate the compounds of formula (I) in the form of nanoparticles which have a surface modifier adsorbed on the surface thereof in an amount sufficient to maintain an effective average particle size of less than 1000 nm.

- 5 Useful surface modifiers are believed to include those which physically adhere to the surface of the compound of formula (I) but do not chemically bond to said compound.

- Suitable surface modifiers can preferably be selected from known organic and inorganic pharmaceutical excipients. Such excipients include various polymers, low molecular weight oligomers, natural products and surfactants. Preferred surface modifiers include nonionic and anionic surfactants.
- 10

- Yet another interesting way of formulating the compounds of formula (I) involves a pharmaceutical composition whereby the compounds of formula (I) are incorporated in hydrophilic polymers and applying this mixture as a coat film over many small beads, thus yielding a composition which can conveniently be manufactured and which is suitable for preparing pharmaceutical dosage forms for oral administration.
- 15

- Said beads comprise a central, rounded or spherical core, a coating film of a hydrophilic polymer and a compound of formula (I) and optionally a seal-coating layer.
- 20

- Materials suitable for use as cores in the beads are manifold, provided that said materials are pharmaceutically acceptable and have appropriate dimensions and firmness. Examples of such materials are polymers, inorganic substances, organic substances, and saccharides and derivatives thereof.
- 25

- It is especially advantageous to formulate the aforementioned pharmaceutical compositions in unit dosage form for ease of administration and uniformity of dosage. Unit dosage form as used herein refers to physically discrete units suitable as unitary dosages, each unit containing a predetermined quantity of active ingredient calculated to produce the desired therapeutic effect in association with the required pharmaceutical carrier. Examples of such unit dosage forms are tablets (including scored or coated tablets), capsules, pills, powder packets, wafers, suppositories, injectable solutions or suspensions and the like, and segregated multiples thereof.
- 30

- Those of skill in the treatment of HIV-infection could determine the effective daily amount from the test results presented here. In general it is contemplated that an effective daily amount would be from 0.01 mg/kg to 50 mg/kg body weight, more preferably from 0.1 mg/kg to 10 mg/kg body weight. It may be appropriate to
- 35

administer the required dose as two, three, four or more sub-doses at appropriate intervals throughout the day. Said sub-doses may be formulated as unit dosage forms, for example, containing 1 to 1000 mg, and in particular 5 to 200 mg of active ingredient per unit dosage form.

5

The exact dosage and frequency of administration depends on the particular compound of formula (I) used, the particular condition being treated, the severity of the condition being treated, the age, weight and general physical condition of the particular patient as well as other medication the individual may be taking, as is well known to those skilled in the art. Furthermore, it is evident that said effective daily amount may be lowered or increased depending on the response of the treated subject and/or depending on the evaluation of the physician prescribing the compounds of the instant invention. The effective daily amount ranges mentioned hereinabove are therefore only guidelines and are not intended to limit the scope or use of the invention to any extent.

15

The present compounds of formula (I) can be used alone or in combination with other therapeutic agents, such as anti-virals, antibiotics, immunomodulators or vaccines for the treatment of viral infections. They may also be used alone or in combination with other prophylactic agents for the prevention of viral infections. The present compounds may be used in vaccines and methods for protecting individuals against viral infections over an extended period of time. The compounds may be employed in such vaccines either alone or together with other compounds of this invention or together with other anti-viral agents in a manner consistent with the conventional utilization of reverse transcriptase inhibitors in vaccines. Thus, the present compounds may be combined with pharmaceutically acceptable adjuvants conventionally employed in vaccines and administered in prophylactically effective amounts to protect individuals over an extended period of time against HIV infection.

20

25

Also, the combination of an antiretroviral compound and a compound of the present invention can be used as a medicine. Thus, the present invention also relates to a product containing (a) a compound of the present invention, and (b) another antiretroviral compound, as a combined preparation for simultaneous, separate or sequential use in the prevention or treatment of retroviral infections, in particular, in the treatment of infections with multi-drug resistant retroviruses. Thus, to combat, prevent or treat HIV infections, or the infection and disease associated with HIV infections, such as Acquired Immunodeficiency Syndrome (AIDS) or AIDS Related Complex (ARC), the compounds of this invention may be co-administered in combination with for instance, binding inhibitors, such as, for example, dextran sulfate, suramine, polyanions, soluble CD4, PRO-542, BMS-806; fusion inhibitors, such as, for example, T20, T1249, RPR

30

35

-37-

- 103611, YK-FH312, IC 9564, 5-helix, D-peptide ADS-J1; co-receptor binding inhibitors, such as, for example, AMD 3100, AMD-3465, AMD7049, AMD3451 (Bicyclams), TAK 779, T-22, ALX40-4C; SHC-C (SCH351125), SHC-D, PRO-140, RPR103611; RT inhibitors, such as, for example, foscarnet and prodrugs; nucleoside
- 5 RTIs, such as, for example, AZT, 3TC, DDC, DDI, D4T, Abacavir, FTC, DAPD (Amdoxovir), dOTC (BCH-10652), fozivudine, DPC 817; nucleotide RTIs, such as, for example, PMEA, PMPA (tenofovir); NNRTIs, such as, for example, nevirapine, delavirdine, efavirenz, 8 and 9-CI TIBO (tivrapipe), loviride, TMC-125, dapivirine, MKC-442, UC 781, UC 782, Capravirine, QM96521, GW420867X, DPC 961,
- 10 DPC963, DPC082, DPC083, TMC-125, calanolide A, SJ-3366, TSAO, 4"-deaminated TSAO, MV150, MV026048, PNU-142721; RNase H inhibitors, such as, for example, SP1093V, PD126338; TAT inhibitors, such as, for example, RO-5-3335, K12, K37; integrase inhibitors, such as, for example, L 708906, L 731988, S-1360; protease inhibitors, such as, for example, amprenavir and prodrug GW908 (fosamprenavir),
- 15 ritonavir, nelfinavir, saquinavir, indinavir, lopinavir, palinavir, BMS 186316, atazanavir, DPC 681, DPC 684, tipranavir, AG1776, mozenavir, DMP-323, GS3333, KNI-413, KNI-272, L754394, L756425, LG-71350, PD161374, PD173606, PD177298, PD178390, PD178392, PNU 140135, TMC-114, maslinic acid, U-140690; glycosylation inhibitors, such as, for example, castanospermine, deoxynojirimycine;
- 20 entry inhibitors CGP64222.

- By administering the compounds of the present invention with other anti-viral agents which target different events in the viral life cycle, the therapeutic effect of these compounds can be potentiated. Combination therapies as described above exert a
- 25 synergistic effect in inhibiting HIV replication because each component of the combination acts on a different site of HIV replication. The use of such combinations may reduce the dosage of a given conventional anti-retroviral agent which would be required for a desired therapeutic or prophylactic effect as compared to when that agent is administered as a monotherapy. These combinations may reduce or eliminate the side
- 30 effects of conventional single anti-retroviral therapy while not interfering with the anti-viral activity of the agents. These combinations reduce potential of resistance to single agent therapies, while minimizing any associated toxicity. These combinations may also increase the efficacy of the conventional agent without increasing the associated toxicity.
- 35 The compounds of the present invention may also be administered in combination with immunomodulating agents, e.g. levamisole, bropirimine, anti-human alpha interferon antibody, interferon alpha, interleukin 2, methionine enkephalin, diethyldithiocarbamate, tumor necrosis factor, naltrexone and the like; antibiotics, e.g. pentamidine isethionate

-38-

and the like; cholinergic agents, e.g. tacrine, rivastigmine, donepezil, galantamine and the like; NMDA channel blockers, e.g. memantine to prevent or combat infection and diseases or symptoms of diseases associated with HIV infections, such as AIDS and ARC, e.g. dementia. A compound of formula (I) can also be combined with another
5 compound of formula (I).

Although the present invention focuses on the use of the present compounds for preventing or treating HIV infections, the present compounds may also be used as inhibitory agents for other viruses, which depend on similar reverse transcriptases for
10 obligatory events in their life cycle.

Examples



The following examples are intended to illustrate the present invention.

Hereinafter, "DMF" is defined as *N,N*-dimethylformamide, "DIPE" is defined as
15 diisopropyl ether, "THF" is defined as tetrahydrofuran, "DMSO" is defined as dimethylsulfoxide, "EtOAc" is defined as ethylacetate.

Examples 1-12 : Synthesis of compounds 1.10, 1.11, 1.18 and 1.9.

20 Scheme 1

-39-

**Example 1. Preparation of intermediate 1b**

N-Chlorosuccinimide (0.025 mol) was added portionwise to a mixture of 5-hydroxy-1-indanone **1a** (0.022 mol) in acetonitrile (60ml). The mixture was stirred and refluxed overnight. H₂O was added and the mixture was extracted with CH₂Cl₂. The organic layer was separated, dried with MgSO₄, filtered, and evaporated. The residue (6g) was purified by column chromatography over silica gel (eluent: cyclohexane/EtOAc 60/40; 15-40μm). Two fractions were collected and evaporated, yielding: 2.2g F1 and 1.3g of starting material. F1 was crystallized from di-isopropyl ether. The precipitate was filtered off and dried yielding 0.9g of intermediate **1b** (22%) (Melting point : 212°C)

Example 2. Preparation of compound 1.10

Sodium hydride (60% in oil) (0.0191 mol) was added portionwise to a mixture of intermediate **1a** (0.0183 mol) in 1,4-dioxane (25ml). The mixture was stirred at room

-40-

temperature for 10 minutes after which 1-methylpyrrolidinone (25ml) was added slowly. Intermediate I-1 (0.0183 mol) was added and the mixture was stirred at 150°C overnight, poured out on ice. The precipitate was filtered, washed with diethyl ether and dried, yielding 4.43g of compound 1.10 (85%). (Melting point: >260°C; (MH⁺) : 343)

5 Example 3. Preparation of compound 1.11

Sodium hydride (60% in oil) (0.0054 mol) was added to a mixture of 1b (0.0049 mol) in 1,4-dioxane (10ml) and stirred for 10 minutes. Subsequently, 1-methylpyrrolidinone (10ml) was added and the mixture was stirred for 10 minutes. Intermediate I-1 (0.0049 mol) was added and the mixture was stirred at 140°C overnight. H₂O was added to the mixture and the mixture was extracted with CH₂Cl₂. The organic layer was washed with K₂CO₃ 10%, dried (MgSO₄), filtered and evaporated, yielding 1.6g of intermediate product. This fraction was crystallized from CH₃CN. The precipitate was filtered off and dried, yielding: 0.46g of compound 1.11 (29%). (Melting point : >260°C; (MH⁺) : 388)

15 Compounds of formula 1.14 to 1.17 are prepared according to the same procedure.

Example 4. Preparation of compound 1.1 (E/Z = 50/50)

Potassium ter-butoxide (0.0018 mol) was added at 0°C to a mixture of diethyl cyanomethylphosphonate (0.0011 mol) in THF (40ml). The mixture was stirred at room temperature for 1 hour. A mixture of compound 1.10 (0.0011 mol) in THF (40ml) was added slowly. The mixture was stirred at room temperature overnight, poured out on ice and extracted with EtOAc. The organic layer was separated, dried (MgSO₄), filtered, and evaporated. The residue (0.48g) was purified by column chromatography over silica gel (eluent: CH₂Cl₂/EtOAc 98/3; 10µm). The pure fractions were collected and the solvent was evaporated, yielding 0.1g product. This fraction was crystallized from di-isopropyl ether. The precipitate was filtered off and dried, yielding 0.086g of compound 1.1 (19%). (Melting point : 225°C, (MH⁺): 366)

25 Compounds of formula 1.2, 1.3, 1.6, 1.7 and 1.8 can be prepared according to the same procedure.

30 Example 5. Preparation of compound 1.18 (E/Z = 70/30)


Potassium ter-butoxide (0.0009 mol) was added portionwise at 5°C to a mixture of diethyl(1-cyanoethyl)phosphonate (0.0009 mol) in THF (8ml) under N₂ flow. The mixture was stirred at 10°C for 30 minutes, then at room temperature for 30 minutes. A solution of compound 1.11 (0.0006 mol) in THF (8ml) was added. The mixture was stirred at room temperature overnight. H₂O was added and the mixture was extracted with CH₂Cl₂. The organic layer was separated, dried (MgSO₄), filtered, and evaporated. The residue (0.25g) was purified by column chromatography over kromasyl (eluent:

35

-41-

CH₂Cl₂/cyclohexane 50/50; 10µm). Two fractions were collected and evaporated yielding: 0.084g F1 (secondary product) and 0.042g of compound 1.18 (18%). (Melting point: 245°C; (MH⁺): 414)

5 Example 6. Preparation of compound 1.9 (E/Z = 85/15)



A mixture of compound 1.10 (0.0005 mol) and *O*-benzylhydroxylamine hydrochloride (0.0008 mol) in EtOH (20ml) was stirred and refluxed for 2 hours. K₂CO₃ 10% was added. The mixture was extracted with EtOAc. The organic layer was separated, dried with MgSO₄, filtered, and evaporated. The residue (0.03g) was crystallized from diethyl ether/ diisopropyl ether. The precipitate was filtered off and dried, yielding 0.03g of compound 1.9 (12%). (Melting point : 134°C)

15 Example 7. Preparation of compound 2.1

A mixture of 6-amino-indan-1-one (0.0003 mol) and intermediate I-1 (0.0003 mol) in HCl 3N (2ml) was stirred and refluxed for 2 hours. The precipitate was filtered, washed with H₂O and diisopropyl ether and dried. The yield of this procedure was 0.06g (52%). This fraction was crystallized from CH₃CN/diisopropyl ether. The precipitate was filtered off and dried, yielding 0.035g of compound 2.1 (30%). (Melting point: >260°C)

20 Example 8. Preparation of compound 2.2 (100 % E)


Potassium ter-butoxide (0.0016 mol) was added at 5°C to a mixture of diethyl cyanomethylphosphonate (0.0016 mol) in THF (3ml) under N₂ flow. The mixture was stirred at room temperature for 1 hour. A solution of compound 2.1 (0.0011 mol) in THF (3ml) was added. The mixture was stirred at room temperature for 2 hours. H₂O was added. The mixture was extracted with CH₂Cl₂. The organic layer was separated, dried (MgSO₄), filtered, and evaporated. The residue (0.82g) was purified by column chromatography over silica gel (15-35µm). The pure fractions were collected and the solvent was evaporated yielding 0.122g product (30%). This fraction was crystallized

-42-

from CH₃CN/ diisopropyl ether. The precipitate was filtered off and dried yielding 0.035g of compound 2.2 (9%). (Melting point: >270°C, (MH⁺): 375)

Scheme 2

5

**Example 9. Preparation of intermediate 3b**

10 Fuming nitric acid (0.362 mol) was added at -40°C to 5-chloroindan-1-one (Int 3a, 26.7 mmol). The mixture was stirred for two hours at -40°C . It was then poured onto ice and extracted with dichloromethane. The organic layer was separated, washed with brine, dried with MgSO_4 , filtered and evaporated. The residue was purified by column chromatography over silica gel (eluent : cyclohexane/AcOEt 65/35; 15-40 μm). The pure fractions were collected and the solvent evaporated yielding 4,15 g of intermediate 3b (73 %). (Melting Point : 129°C)

Example 10. Preparation of intermediate 3c

20 In a Parr hydrogenation apparatus, 0.5 g of Raney Nickel was added to a solution of intermediate 3b (8.60 mmol) in a mixture of THF and MeOH (6/1). The vessel was flushed with nitrogen and put under an hydrogen atmosphere (3 bars). The mixture was

-43-

stirred for one hour at room temperature, filtered over celite and evaporated to dryness to yield 1.50 g of intermediate 3c (96 %). (Melting Point : 214°C)

Example 11. Preparation of intermediate 3d (E/Z=89/11)

- 5 Potassium *tert*-butoxide (56.4 mmol) was added portionwise at 0°C to a solution of cyanomethylphosphonate (56.4 mmol) in THF. The mixture was stirred for 15 min at 15°C. Then a solution of intermediate 3c (14.1 mmol) in THF was added dropwise at 0°C. The reaction mixture was stirred at room temperature overnight, poured onto water, acidified with 3M hydrochloric acid and extracted with dichloromethane. The organic layer was separated, washed with a 10% solution of potassium carbonate, with brine, dried over magnesium sulfate, filtered and evaporated. The residue was purified by column chromatography over silica gel (eluent : dichloromethane; 35-70 µm). The pure fractions were collected and the solvent evaporated yielding 1.88 g of intermediate 3d (65 %). (Melting Point : 196°C)

15


Example 12. Preparation of compound 2.3 (Z 100 %)

- Intermediates 3d and I-2 were intimately ground together and fused with a heating gun. The residue was taken up with a 90/10 mixture of dichloromethane and methanol and with a 10% solution of potassium carbonate. The organic layer was separated, washed with a brine solution, dried over magnesium sulfate, filtered and evaporated. The residue was purified by column chromatography over silica gel (eluent : dichloromethane/AcOEt 85/15; 15-40 µm). Two fractions were collected and evaporated yielding 0.126g of compound 2.3 (13 %) and 0.104g of another isomer (11 %). Each fraction was recrystallized in acetonitrile, yielding compound 2.3 (Melting Point : 248-249°C) and its isomer (Melting point >250°C).

25

Examples 13-14 : Synthesis of compound 1.20

Scheme B



30


Example 13. Preparation of compound 1.19

A solution of 4,6-dimethyl-5-hydroxyindan-1-one **4a** (2.07 mmol) in dioxane and NMP was cooled to 0°C. Sodium hydride (2.28 mmol) was added and the mixture was stirred at room temperature for 30 min. Then a solution of intermediate **2** in NMP was added.

- 5 The reaction mixture was refluxed for 7 days and was then evaporated to dryness. The residue was washed with a 90/10 mixture of dichloromethane and methanol to yield 0.601 g of compound **1.19** (78 %).

Example 14. Preparation of compound 1.20

- 10 Potassium *tert*-butoxide (5.26 mmol) was added portionwise at 0°C to a solution of cyanomethylphosphonate (5.26 mmol) in THF. The mixture was stirred for 15 min at 15°C. Then a solution of compound **1.19** (1.75 mmol) in THF was added dropwise at 0°C. The reaction mixture was stirred at room temperature for 2 hours, poured onto 0.5M hydrochloric acid and extracted with a 90/10 mixture of dichloromethane and methanol.
- 15 The organic layer was separated, washed with water and brine, dried over magnesium sulfate, filtered and evaporated. The residue was washed with dichloromethane and diisopropylether yielding 0.205 g of compound **1.20** (30 %). (Melting Point >250°C)

20 Table 1


N°	X ¹	R ^{17a}	R ^{17b}	R ^{17c}	R ^{17d}	R ^{4a}	R ^{4b}	Y	Q
1.1	O	=CH-CN		H	H	H	H	H	H
1.2	O	=CH-CN		H	H	Cl	Cl	H	H
1.3	O	=CH-CN		H	H	Cl	H	H	H
1.4	O	-CH ₂ CN	-H	H	H	H	H	H	H
1.5	O	-CH ₂ CN	-H	H	H	Cl	H	H	H
1.6	O	=CH-CN		H	H	Br	H	H	H
1.7	O	=CH-CN		H	H	CH ₃	H	H	H
1.8	O	=CH-CN		H	H	CH ₃	Cl	H	H

-45-

N°	X ¹	R ^{17a}	R ^{17b}	R ^{17c}	R ^{17d}	R ^{4a}	R ^{4b}	Y	Q
1.9	O	=N-O-CH ₂ - phenyl		H	H	H	H	H	H
1.10	O	=O		H	H	H	H	H	H
1.11	O	=O		H	H	Cl	H	H	H
1.12	NH	=CH-CN		H	H	H	H	H	H
1.13	NH	=CH-CN		H	H	Cl	H	H	H
1.14	O	=O		H	H	Cl	Cl	H	H
1.15	O	=O		H	H	Br	H	H	H
1.16	O	=O		H	H	CH ₃	H	H	H
1.17	O	=O		H	H	CH ₃	Cl	H	H
1.18	O	=C(CH ₃)(CN)		H	H	Cl	H	H	H
1.19	O	=O		H	H	CH ₃	CH ₃	H	H
1.20	O	=N-CN		H	H	CH ₃	CH ₃	H	H

For compounds 1.1-1.3 and 1.6-1.18, R^{17a} and R^{17b} taken together form a double bond.

Table 2




5

N°	X ₁	R ^{17a}	R ^{17b}	R ^{17c}	R ^{17d}	R ^{4a}	R ^{4b}	Y	Q
2.1	NH	=O		H	H	H	H	H	H
2.2	NH	=CH-CN		H	H	H	H	H	H
2.3	NH	=CH-CN		H	H	Cl	H	H	H
2.4	NH	=CH-CN		H	H	CH ₃	CH ₃	H	H
2.5	O	=CH-CN		H	H	CH ₃	CH ₃	H	H
2.6	O	=CH-CN		H	H	Cl	H	H	H

For compounds 2.1 – 2.30, R^{17a} and R^{17b} taken together form a double bond.

Table 3




No.	X ¹	Q	R ⁴	R ^{17a}	R ^{17b}	R ^{2a}	R ^{2b}
3.1	O	H	Cl	=CH-CN		-CH ₂ -CN	H
3.2	O	H	Cl	=CH-CN		-NO ₂	H
3.3	O	H	Cl	=CH-CN			H
3.4	O	H	Cl	=CH-CN		-CO-NH ₂	H
3.5	O	H	Cl	=CH-CN		F	H
3.6	O	H	Cl	=CH-CN		Cl	H
3.7	O	H	Cl	=CH-CN		Br	H
3.8	O	H	Cl	=CH-CN		CN	OH
3.9	O	H	Cl	=CH-CN		CN	Cl
3.10	O	H	Cl	=CH-CN		F	F
3.11	O	H	Cl	=CH-CN		-CH ₂ -CO-NH ₂	H
3.12	O	CN	Cl	=CH-CN		CN	H
3.13	NH	CN	Cl	=CH-CN		CN	H
3.14	NH	H	Cl	=CH-CN		-CH ₂ -CN	H
3.15	NH	H	Cl	=CH-CN		-NO ₂	H
3.16	NH	H	Cl	=CH-CN			H
3.17	NH	H	Cl	=CH-CN		-CO-NH ₂	H
3.18	NH	H	Cl	=CH-CN		F	H
3.19	NH	H	Cl	=CH-CN		Cl	H
3.20	NH	H	Cl	=CH-CN		Br	H
3.21	NH	H	Cl	=CH-CN		CN	OH
3.22	NH	H	Cl	=CH-CN		CN	Cl
3.23	NH	H	Cl	=CH-CN		F	F
3.24	NH	H	Cl	=CH-CN		-CH ₂ -CO-NH ₂	H
3.25	O	H	Cl	=CH-CO-NH ₂		CN	H

-47-

No.	X ¹	Q	R ⁴	R ^{17a}	R ^{17b}	R ^{2a}	R ^{2b}
3.26	NH	H	Cl	=CH-CO-NH ₂		CN	H
3.27	O	H	Cl	-CH ₂ -CO-NH ₂	H	CN	H
3.28	NH	H	Cl	-CH ₂ -CO-NH ₂	H	CN	H
3.29	O	H	Cl	-CH=CH-CN	H	CN	H
3.30	NH	H	Cl	-CH=CH-CN	H	CN	H

Table 4



5

No.	X ¹	Q	R ^{4a}	R ^{4b}	R ^{17a}	R ^{17b}	R ^{2a}	R ^{2b}
4.1	O	H	CH ₃	CH ₃	=CH-CN		-CH ₂ -CN	H
4.2	O	H	CH ₃	CH ₃	=CH-CN		-NO ₂	H
4.3	O	H	CH ₃	CH ₃	=CH-CN			H
4.4	O	H	CH ₃	CH ₃	=CH-CN		-CO-NH ₂	H
4.5	O	H	CH ₃	CH ₃	=CH-CN		F	H
4.6	O	H	CH ₃	CH ₃	=CH-CN		Cl	H
4.7	O	H	CH ₃	CH ₃	=CH-CN		Br	H
4.8	O	H	CH ₃	CH ₃	=CH-CN		CN	OH
4.9	O	H	CH ₃	CH ₃	=CH-CN		CN	Cl
4.10	O	H	CH ₃	CH ₃	=CH-CN		F	F
4.11	O	H	CH ₃	CH ₃	=CH-CN		-CH ₂ -CO-NH ₂	H
4.12	NH	CN	CH ₃	CH ₃	=CH-CN		CN	H
4.13	O	CN	CH ₃	CH ₃	=CH-CN		CN	H
4.14	O	-CO-N(CH ₃) ₂	CH ₃	CH ₃	=CH-CN		CN	H
4.15	O	-CH=CH ₂	CH ₃	CH ₃	=CH-CN		CN	H
4.16	O	-CO-NH ₂	CH ₃	CH ₃	=CH-CN		CN	H
4.17	O	-C≡CH	CH ₃	CH ₃	=CH-CN		CN	H
4.18	O	-COOCH ₃	CH ₃	CH ₃	=CH-CN		CN	H
4.19	O	H	CH ₃	CH ₃	-CH ₂ -CONH ₂	H	CN	H

No.	X ¹	Q	R ^{4a}	R ^{4b}	R ^{17a}	R ^{17b}	R ^{2a}	R ^{2b}
4.20	NH	H	CH ₃	CH ₃	-CH ₂ -CONH ₂	H	CN	H
4.21	O	H	CH ₃	CH ₃	=CH-CONH ₂		CN	H
4.22	NH	H	CH ₃	CH ₃	-CH=CH-CN	H	CN	H
4.23	O	H	CH ₃	CH ₃	-CH=CH-CN	H	CN	H

Formulation examples

Capsules

Active ingredient, *in casu* a compound of formula (I), can be dissolved in organic solvent such as ethanol, methanol or methylene chloride, preferably, a mixture of ethanol and methylene chloride. Polymers such as polyvinylpyrrolidone copolymer with vinyl acetate (PVP-VA) or hydroxypropylmethylcellulose (HPMC), typically 5 mPa.s, can be dissolved in organic solvents such as ethanol, methanol methylene chloride. Suitably the polymer can be dissolved in ethanol. The polymer and compound solutions can be mixed and subsequently spray dried. The ratio of compound/polymer can be selected from 1/1 to 1/6. Intermediate ranges can be 1/1.5 and 1/3. A suitable ratio can be 1/6. The spray-dried powder, a solid dispersion, can subsequently be filled in capsules for administration. The drug load in one capsule can range between 50 and 100 mg depending on the capsule size used.

Film-coated Tablets

Preparation of Tablet Core

A mixture of 100 g of active ingredient, *in casu* a compound of formula (I), 570 g lactose and 200 g starch can be mixed well and thereafter humidified with a solution of 5 g sodium dodecyl sulfate and 10 g polyvinylpyrrolidone in about 200 ml of water. The wet powder mixture can be sieved, dried and sieved again. Then there can be added 100 g microcrystalline cellulose and 15 g hydrogenated vegetable oil. The whole can be mixed well and compressed into tablets, giving 10.000 tablets, each comprising 10 mg of the active ingredient.

Coating

To a solution of 10 g methylcellulose in 75 ml of denaturated ethanol there can be added a solution of 5 g of ethylcellulose in 150 ml of dichloromethane. Then there can be added 75 ml of dichloromethane and 2.5 ml 1,2,3-propanetriol. 10 g of polyethylene glycol can be molten and dissolved in 75 ml of dichloromethane. The latter solution can be added to the former and then there can be added 2.5 g of magnesium octadecanoate, 5 g of polyvinylpyrrolidone and 30 ml of concentrated color suspension and the whole can be homogenated. The tablet cores can be coated with the thus obtained mixture in a coating apparatus.

Antiviral analyses:

The compounds of the present invention were examined for anti-viral activity in a cellular assay. The assay demonstrated that these compounds exhibited potent anti-HIV activity against a wild type laboratory HIV strain (HIV-1 strain LAI). The cellular assay was performed according to the following procedure.

Cellular Assay Experimental Method:

HIV- or mock-infected MT4 cells were incubated for five days in the presence of various concentrations of the inhibitor. At the end of the incubation period, all HIV-infected cells have been killed by the replicating virus in the control cultures in the absence of any inhibitor. Cell viability is measured by measuring the concentration of MTT, a yellow, water soluble tetrazolium dye that is converted to a purple, water insoluble formazan in the mitochondria of living cells only. Upon solubilization of the resulting formazan crystals with isopropanol, the absorbance of the solution is monitored at 540nm. The values correlate directly to the number of living cells remaining in the culture at the completion of the five day incubation. The inhibitory activity of the compound was monitored on the virus-infected cells and was expressed as EC_{50} and EC_{90} . These values represent the amount of the compound required to protect 50% and 90%, respectively, of the cells from the cytopathogenic effect of the virus. The toxicity of the compound was measured on the mock-infected cells and was expressed as CC_{50} , which represents the concentration of compound required to inhibit the growth of the cells by 50%. The selectivity index (SI) (ratio CC_{50}/EC_{50}) is an indication of the selectivity of the anti-HIV activity of the inhibitor. Wherever results are reported as e.g. pEC_{50} or pCC_{50} values, the result is expressed as the negative logarithm of the result expressed as EC_{50} or CC_{50} respectively.

Antiviral spectrum:

Because of the increasing emergence of drug resistant HIV strains, the present compounds were tested for their potency against clinically isolated HIV strains harboring several mutations. These mutations are associated with resistance to reverse transcriptase inhibitors and result in viruses that show various degrees of phenotypic cross-resistance to the currently commercially available drugs such as for instance AZT and delavirdine.

35

The antiviral activity of the compound of the present invention has been evaluated in the presence of wild type HIV and HIV mutants bearing mutations at the reverse transcriptase gene. The activity of the compounds is evaluated using a cellular assay and the residual activity is expressed in pEC_{50} values. Column A contains the pEC_{50} against

-50-

- strain A (Strain A contains mutation 100I in HIV reverse transcriptase), Column B contains the pEC_{50} against strain B (Strain B contains mutation 100I and 103N in HIV reverse transcriptase), Column C contains the pEC_{50} against strain C (Strain C contains mutation 103N in HIV reverse transcriptase), Column D contains the pEC_{50} against strain D (Strain D contains mutation 181C in HIV reverse transcriptase), Column E contains the pEC_{50} against strain E (Strain E contains mutation 188L in HIV reverse transcriptase), Column F contains the pEC_{50} against strain F (Strain F contains mutation 227C in HIV reverse transcriptase), and Column G contains the pEC_{50} against strain G (Strain G contains mutation 106A and 227L in HIV reverse transcriptase),). Column
- 5 IIIB displays the pEC_{50} value against wild type HIV-LAI strain. ND, not determined.
- 10


Table 5

Compound number	IIIB	A	B	C	D	E	F	G
1.1	8.7	6.3	5	7.7	6.5	5.7	6.6	7.0
1.18	8.6	6.7	6.1	8.0	6.8	6.3	6.7	7.3
1.19	8.0	5.7	5.6	6.0	5.5	5.2	6.0	6.5

-51-

Claims

1. A compound of formula (I)

a *N*-oxide, a pharmaceutically acceptable addition salt, a quaternary amine or a


5 stereochemically isomeric form thereof, wherein

 $-a^1=a^2-a^3=a^4-$ represents a bivalent radical of formula $-\text{CH}=\text{CH}-\text{CH}=\text{CH}-$ (a-1); $-\text{N}=\text{CH}-\text{CH}=\text{CH}-$ (a-2); $-\text{N}=\text{CH}-\text{N}=\text{CH}-$ (a-3);10 $-\text{N}=\text{CH}-\text{CH}=\text{N}-$ (a-4); $-\text{N}=\text{N}-\text{CH}=\text{CH}-$ (a-5); $-b^1-b^2-b^3-$ represents a bivalent radical of formula $-\text{CH}_2-\text{CH}_2-\text{CH}_2-$ (b-1);n is 0, 1, 2, 3 or 4; and in case $-a^1=a^2-a^3=a^4-$ is (a-1), then n may also be 5;

15 m is 0, 1, 2, 3;

q is 0, 1 or 2;

p is 1 or 2;

 R^1 is hydrogen; aryl; formyl; C_{1-6} alkylcarbonyl; C_{1-6} alkyl; C_{1-6} alkyloxycarbonyl; C_{1-6} alkyl substituted with formyl, C_{1-6} alkylcarbonyl, C_{1-6} alkyloxycarbonyl,20 C_{1-6} alkylcarbonyloxy; C_{1-6} alkyloxy C_{1-6} alkylcarbonyl substituted with C_{1-6} alkyloxycarbonyl;each R^2 independently is hydroxy, halo, C_{1-6} alkyl optionally substituted with cyano or with $-\text{C}(=\text{O})\text{R}^6$, C_{3-7} cycloalkyl, C_{2-6} alkenyl optionally substituted with one or morehalogen atoms or cyano, C_{2-6} alkynyl optionally substituted with one or more halogen25 atoms or cyano, C_{1-6} alkyloxycarbonyl, carboxyl, cyano, nitro, $\text{NR}^{13}\text{R}^{14}$,polyhalomethyl, polyhalomethylthio, $-\text{S}(=\text{O})_p\text{R}^6$, $-\text{NH}-\text{S}(=\text{O})_p\text{R}^6$, $-\text{C}(=\text{O})\text{R}^6$, $-\text{NHC}(=\text{O})\text{H}$, $-\text{C}(=\text{O})\text{NHNH}_2$, $-\text{NHC}(=\text{O})\text{R}^6$, $-\text{C}(=\text{NH})\text{R}^6$ or a radical of formulawherein each A_1 independently is N, CH or CR^6 ; and30 A_2 is NH, O, S or NR^6 ; X_1 is $-\text{NR}^5-$, $-\text{NH}-\text{NH}-$, $-\text{N}=\text{N}-$, $-\text{O}-$, $-\text{C}(=\text{O})-$, C_{1-4} alkanediyl, $-\text{CHOH}-$, $-\text{S}-$, $-\text{S}(=\text{O})_p-$, $-\text{NR}^{13}-\text{C}(=\text{O})-$, $-\text{C}(=\text{O})-\text{NR}^{13}-$, $-\text{X}_2-\text{C}_{1-4}$ alkanediyl- or $-\text{C}_{1-4}$ alkanediyl- X_2- ;

-52-

X_2 is $-NR^5$ -, $-NH-NH$ -, $-N=N$ -, $-O$ -, $-C(=O)$ -, $-CHOH$ -, $-S$ -, $-S(=O)_p$ -;

R^3 is hydrogen, halo, C_{1-6} alkyl, $NR^{13}R^{14}$ -, $-C(=O)-NR^{13}R^{14}$ -, $-C(=O)-R^{15}$ -, $-CH=N-NH-C(=O)-R^{16}$ -, $-C(=N-O-R^8)-C_{1-4}$ alkyl, R^7 or $-X_3-R^7$; or C_{1-6} alkyl substituted with one or more substituents each independently selected from halo, hydroxy, cyano, NR^9R^{10} -, $-C(=O)-NR^9R^{10}$ -, $-C(=O)-C_{1-6}$ alkyl or R^7 , and in addition to said list of substituents, two geminal hydrogen atoms of said C_{1-6} alkyl may also be replaced by a C_{2-5} alkanediyl thus forming a spiro ring; C_{1-6} alkoxy C_{1-6} alkyl optionally substituted with one or more substituents each independently selected from hydroxy, cyano, NR^9R^{10} -, $-C(=O)-NR^9R^{10}$ -, $-C(=O)-C_{1-6}$ alkyl or R^7 ; C_{2-6} alkenyl substituted with one or more substituents each independently selected from halo, hydroxy, cyano, NR^9R^{10} -, $-C(=O)-NR^9R^{10}$ -, $-C(=O)-C_{1-6}$ alkyl or R^7 ; C_{2-6} alkynyl substituted with one or more substituents each independently selected from halo, hydroxy, cyano, NR^9R^{10} -, $-C(=O)-NR^9R^{10}$ -, $-C(=O)-C_{1-6}$ alkyl or R^7 ;

X_3 is $-NR^5$ -, $-NH-NH$ -, $-N=N$ -, $-O$ -, $-C(=O)$ -, $-S$ -, $-S(=O)_p$ -, $-X_2-C_{1-4}$ alkanediyl-, $-C_{1-4}$ alkanediyl- X_{2a} -, $-C_{1-4}$ alkanediyl- $X_{2b}-C_{1-4}$ alkanediyl-, $-C(=N-OR^8)-C_{1-4}$ alkanediyl-; with X_{2a} being $-NH-NH$ -, $-N=N$ -, $-O$ -, $-C(=O)$ -, $-S$ -, $-S(=O)_p$ -; and with X_{2b} being $-NH-NH$ -, $-N=N$ -, $-C(=O)$ -, $-S$ -, $-S(=O)_p$ -;

R^4 is halo, hydroxy, C_{1-6} alkyl, C_{2-6} alkenyl, C_{2-6} alkynyl, C_{3-7} cycloalkyl, C_{1-6} alkyloxy, cyano, nitro, polyhalo C_{1-6} alkyl, polyhalo C_{1-6} alkyloxy, $-C(=O)-NR^{13}R^{14}$ -, C_{1-6} alkyloxycarbonyl, C_{1-6} alkylcarbonyl, formyl, $-NR^{13}R^{14}$ or R^7 ;

R^5 is hydrogen, aryl, formyl; C_{1-6} alkylcarbonyl; C_{1-6} alkyl; C_{1-6} alkyloxycarbonyl; C_{1-6} alkyl substituted with formyl, C_{1-6} alkylcarbonyl, C_{1-6} alkyloxycarbonyl or C_{1-6} alkylcarbonyloxy; C_{1-6} alkoxy C_{1-6} alkylcarbonyl substituted with C_{1-6} alkyloxycarbonyl;

R^6 is C_{1-4} alkyl, $NR^{13}R^{14}$ or polyhalo C_{1-4} alkyl;

R^7 is a monocyclic, bicyclic or tricyclic saturated, partially saturated or aromatic carbocycle or a monocyclic, bicyclic or tricyclic saturated, partially saturated or aromatic heterocycle, wherein each of said carbocyclic or heterocyclic ring systems may optionally be substituted with one, two, three, four or five substituents each independently selected from halo, hydroxy, mercapto, C_{1-6} alkyl, hydroxy C_{1-6} alkyl, amino C_{1-6} alkyl, mono or di(C_{1-6} alkyl)amino C_{1-6} alkyl, formyl, C_{1-6} alkylcarbonyl, C_{3-7} cycloalkyl, C_{1-6} alkyloxy, C_{1-6} alkyloxycarbonyl, C_{1-6} alkylthio, cyano, nitro, polyhalo C_{1-6} alkyl, polyhalo C_{1-6} alkyloxy, aminocarbonyl, $-CH(=N-O-R^8)$, R^{7a} -, $-X_3-R^{7a}$ or $R^{7a}-C_{1-4}$ alkyl;

R^{7a} is a monocyclic, bicyclic or tricyclic saturated, partially saturated or aromatic carbocycle or a monocyclic, bicyclic or tricyclic saturated, partially saturated or aromatic heterocycle, wherein each of said carbocyclic or heterocyclic ring systems

- may optionally be substituted with one, two, three, four or five substituents each independently selected from halo, hydroxy, mercapto, C₁₋₆alkyl, hydroxyC₁₋₆alkyl, aminoC₁₋₆alkyl, mono or di(C₁₋₆alkyl)aminoC₁₋₆alkyl, formyl, C₁₋₆alkylcarbonyl, C₃₋₇cycloalkyl, C₁₋₆alkyloxy, C₁₋₆alkyloxycarbonyl, C₁₋₆alkylthio, cyano, nitro,
- 5 polyhaloC₁₋₆alkyl, polyhaloC₁₋₆alkyloxy, -C(=O)-NR¹³R¹⁴, -CH(=N-O-R⁸); R⁸ is hydrogen, C₁₋₄alkyl, aryl, arylC₁₋₄alkyl;
- R⁹ and R¹⁰ each independently are hydrogen; hydroxy; C₁₋₆alkyl; C₁₋₆alkyloxy; C₁₋₆alkylcarbonyl; C₁₋₆alkyloxycarbonyl; NR¹³R¹⁴; -C(=O)-NR¹³R¹⁴; -CH(=NR¹¹) or R⁷, wherein each of the aforementioned C₁₋₆alkyl groups may optionally and each
- 10 individually be substituted with one or two substituents each independently selected from hydroxy, C₁₋₆alkyloxy, hydroxyC₁₋₆alkyloxy, carboxyl, C₁₋₆alkyloxycarbonyl, cyano, imino, NR¹³R¹⁴, polyhalomethyl, polyhalomethyloxy, polyhalomethylthio, -S(=O)_pR⁶, -NH-S(=O)_pR⁶, -C(=O)R⁶, -NHC(=O)H, -C(=O)NHNH₂, -NHC(=O)R⁶, -C(=NH)R⁶, R⁷; or
- 15 R⁹ and R¹⁰ may be taken together to form a bivalent or trivalent radical of formula
- | | | |
|----|---|-------|
| | -CH ₂ -CH ₂ -CH ₂ -CH ₂ - | (d-1) |
| | -CH ₂ -CH ₂ -CH ₂ -CH ₂ -CH ₂ - | (d-2) |
| | -CH ₂ -CH ₂ -O-CH ₂ -CH ₂ - | (d-3) |
| | -CH ₂ -CH ₂ -S-CH ₂ -CH ₂ - | (d-4) |
| 20 | -CH ₂ -CH ₂ -NR ¹² -CH ₂ -CH ₂ - | (d-5) |
| | -CH ₂ -CH=CH-CH ₂ - | (d-6) |
| | =CH-CH=CH-CH=CH- | (d-7) |
- R¹¹ is cyano; C₁₋₄alkylcarbonyl; C₁₋₄alkyloxycarbonyl; -C(=O)-NR¹³R¹⁴; or C₁₋₄alkyl optionally substituted with C₁₋₄alkyloxy, cyano, NR¹³R¹⁴ or -C(=O)-NR¹³R¹⁴;
- 25 R¹² is hydrogen or C₁₋₄alkyl;
- R¹³ and R¹⁴ each independently are hydrogen, Het, C₁₋₆alkyl optionally substituted with cyano or aminocarbonyl, C₂₋₆alkenyl optionally substituted with cyano or aminocarbonyl, C₂₋₆alkynyl optionally substituted with cyano or aminocarbonyl;
- R¹⁵ is C₁₋₆alkyl optionally substituted with cyano or -C(=O)-NR¹³R¹⁴;
- 30 R¹⁶ is R⁷ or C₁₋₆alkyl optionally substituted with cyano or -C(=O)-NR¹³R¹⁴;
- R¹⁷, if present, each independently is cyano, halo, hydroxy, -C(=O)-NR¹³R¹⁴, C₁₋₆alkyl optionally substituted with one or more substituents independently selected from cyano, -C(=O)-NR¹³R¹⁴ or halo; C₂₋₆alkenyl optionally substituted with one or more substituents independently selected from cyano, -C(=O)-NR¹³R¹⁴ or halo; C₂₋₆alkynyl
- 35 optionally substituted with one or more substituents independently selected from cyano, -C(=O)-NR¹³R¹⁴ or halo; and, where possible, R¹⁷ may also be attached to the -b¹-b²-b³- moiety by a double bond whereby R¹⁷ is then =O, =S, =NH, =N-R¹⁵, =N-

-54-

R^7 , $=N-O-R^{15}$, $=N-O-R^7$, $=CH_2$, $=CH-C(=O)-NR^{13}R^{14}$, $=CH-R^7$, or $=CH-R^{15}$;

wherein $=CH_2$ may optionally be substituted with cyano, hydroxy, halo, nitro;

Q represents hydrogen, C_{1-6} alkyl, halo, polyhalo C_{1-6} alkyl, $-C(=O)-NR^{13}R^{14}$, or $-NR^9R^{10}$;

Z is C-Y, wherein,

- 5 Y represents hydrogen, hydroxy, halo, C_{1-6} alkyl, C_{3-7} cycloalkyl, C_{1-6} alkyloxy, C_{1-6} alkyloxycarbonyl, carbonyl, cyano, nitro, $NR^{13}R^{14}$, polyhalomethyl, polyhalomethoxy, polyhalomethylthio, $-S(=O)_pR^8$, $-NH-S(=O)R^8$, $-NH-SO_2-R^8$, $-NH-SO_2-(C_{1-4}alkanedyl)-CO-N(R^8)_2$, $-C(=O)R^8$, $-NHC(=O)H$, $-C(=O)NHNH_2$, $-NHC(=O)R^8$, $-C(=O)-NH-R^8$, $-C(=NH)R^8$, aryl; or C_{2-6} alkenyl optionally substituted with one or more halo atoms;

10 C_{2-6} alkynyl optionally substituted with one or more halo atoms;

C_{1-6} alkyl substituted with cyano or with $-C(=O)R^8$;


aryl is phenyl or phenyl substituted with one, two, three, four or five substituents each independently selected from halo, hydroxy, mercapto, C_{1-6} alkyl, hydroxy C_{1-6} alkyl, C_{1-6} alkyl $NR^{13}R^{14}$, C_{1-6} alkylcarbonyl, C_{3-7} cycloalkyl, C_{1-6} alkyloxy, C_{1-6} alkyloxycarbonyl, C_{1-6} alkylthio, cyano, nitro, polyhalo C_{1-6} alkyl, polyhalo C_{1-6} alkyloxy, $-C(=O)-NR^{13}R^{14}$, R^7 or $-X_3-R^7$;

Het is a monocyclic, bicyclic or tricyclic saturated, partially saturated or aromatic heterocycle, wherein each of said carbocyclic or heterocyclic ring systems may

- 20 optionally be substituted with one, two, three, four or five substituents each independently selected from halo, hydroxy, mercapto, C_{1-6} alkyl, hydroxy C_{1-6} alkyl, amino C_{1-6} alkyl, mono or di(C_{1-6} alkyl)amino C_{1-6} alkyl, formyl, C_{1-6} alkylcarbonyl, C_{3-7} cycloalkyl, C_{1-6} alkyloxy, C_{1-6} alkyloxycarbonyl, C_{1-6} alkylthio, cyano, nitro, polyhalo C_{1-6} alkyl, polyhalo C_{1-6} alkyloxy, $-C(=O)-NR^{13}R^{14}$, $-CH(=N-O-R^8)$.

25 2. A compound according to claim 1 wherein one or more of the following limitations (a) - (v) apply.

(a) $-a^1=a^2-a^3=a^4-$ represents a bivalent radical of formula



30 (b) n is 0, 1, 2, 3;


(c) m is 0, 1 or 2;

(d) R^1 is hydrogen; formyl; C_{1-6} alkylcarbonyl; C_{1-6} alkyl; C_{1-6} alkyloxycarbonyl; C_{1-6} alkylcarbonyl, C_{1-6} alkyloxycarbonyl;

35 (e) each R^2 independently is hydroxy, halo, C_{1-6} alkyl optionally substituted with cyano or $-C(=O)R^6$, C_{3-7} cycloalkyl, C_{2-6} alkenyl optionally substituted with one or more halogen atoms or cyano, C_{2-6} alkynyl optionally substituted with one or more halogen atoms or cyano, C_{1-6} alkyloxycarbonyl, carboxyl, cyano, nitro, amino, mono(C_{1-6} alkyl)amino, di(C_{1-6} alkyl)amino, polyhalomethyl, polyhalomethylthio,

-55-

-S(=O)_pR⁶, -NH-S(=O)_pR⁶, -C(=O)R⁶, -NHC(=O)H, -C(=O)NHNH₂,
 -NHC(=O)R⁶, -C(=NH)R⁶ or a radical of formula



wherein each A₁ independently is N, CH or CR⁶; and

- 5 A₂ is NH, O, S or NR⁶;
- (f) X₁ is -NR⁵-, -NH-NH-, -N=N-, -O-, -C(=O)-, C₁₋₄alkanediyl, -CHOH-, -S-, -S(=O)_p-,
 -, -NR¹³-C(=O)-, -C(=O)-NR¹³-, -X₂-C₁₋₄alkanediyl- or -C₁₋₄alkanediyl-X₂-;
- (g) X₂ is -NR⁵-, -O-;
- (h) R³ is hydrogen, halo, C₁₋₆alkyl, NR¹³R¹⁴, -C(=O)-NR¹³R¹⁴, -C(=O)-R¹⁵, -X₃-R⁷;
- 10 C₁₋₆alkyl substituted with one or more substituents each independently selected
 from cyano, R⁷ or -C(=O)-NR⁹R¹⁰; C₂₋₆alkenyl substituted with one or more
 substituents each independently selected from halo, cyano or -C(=O)-NR⁹R¹⁰ or
 R⁷; or C₂₋₆alkynyl substituted with one or more substituents each independently
 selected from halo, cyano, -C(=O)-NR⁹R¹⁰ or R⁷;
- 15 (i) X₃ is -NR⁵-, -NH-NH-, -N=N-, -O- or -S-
- (j) R⁴ is halo, hydroxy, C₁₋₆alkyl, C₂₋₆alkenyl, C₂₋₆alkynyl, C₁₋₆alkyloxy, cyano, nitro,
 polyhaloC₁₋₆alkyl, polyhaloC₁₋₆alkyloxy, -C(=O)-NR¹³R¹⁴, C₁₋₆alkyloxycarbonyl,
 C₁₋₆alkylcarbonyl, formyl, -NR¹³R¹⁴ or R⁷;
- (k) R⁵ is hydrogen; formyl; C₁₋₆alkylcarbonyl; C₁₋₆alkyl or C₁₋₆alkyloxycarbonyl;
- 20 (l) R⁶ is C₁₋₄alkyl, NR¹³R¹⁴ or polyhaloC₁₋₄alkyl;
- (m) R⁷ is a monocyclic or bicyclic, partially saturated or aromatic carbocycle or a
 monocyclic or bicyclic, partially saturated or aromatic heterocycle, wherein each of
 said carbocyclic or heterocyclic ring systems may optionally be substituted with one,
 two or three substituents each independently selected from halo, hydroxy, mercapto,
- 25 C₁₋₆alkyl, hydroxyC₁₋₆alkyl, aminoC₁₋₆alkyl, C₁₋₆alkylcarbonyl, C₁₋₆alkyloxy,
 C₁₋₆alkyloxycarbonyl, C₁₋₆alkylthio, cyano, nitro, polyhaloC₁₋₆alkyl,
 polyhaloC₁₋₆alkyloxy or aminocarbonyl;
- (n) R⁸ is hydrogen, C₁₋₄alkyl or aryl(C₁₋₄alkyl);
- (o) R⁹ and R¹⁰ each independently are hydrogen; C₁₋₆alkyl; C₁₋₆alkyloxy;
- 30 C₁₋₆alkylcarbonyl or C₁₋₆alkyloxycarbonyl;
- (p) R¹³ and R¹⁴ each independently are hydrogen or C₁₋₆alkyl;
- (q) R¹⁵ is C₁₋₆alkyl optionally substituted with cyano or -C(=O)-NR¹³R¹⁴;
- (r) R¹⁷ is cyano, halo, hydroxy, -C(=O)-NR¹³R¹⁴, C₁₋₆alkyl optionally substituted with
 cyano, -C(=O)-NR¹³R¹⁴ or halo; C₂₋₆alkenyl optionally substituted with cyano or -
 C(=O)-NR¹³R¹⁴; C₂₋₆alkynyl optionally substituted with cyano or -C(=O)-NR¹³R¹⁴;
- 35 and, where possible, R¹⁷ may also be attached to the -b¹-b²-b³- moiety by a double

-56-

bond whereby R^{17} is then $=O$, $=S$, $=NH$, $=N-R^{15}$, $=N-R^7$, $=N-O-R^{15}$, $=N-O-R^7$, $=CH_2$, $=CH-C(=O)-NR^{13}R^{14}$, $=CH-R^7$, or $=CH-R^{15}$; wherein $=CH_2$ may optionally be substituted with cyano, hydroxy, halo, nitro;

(s) Q represents hydrogen, C_{1-6} alkyl or $-NR^9R^{10}$;

5 (t) Y represents hydrogen, hydroxy, halo, C_{1-6} alkyl, C_{1-6} alkyloxy, cyano, nitro, $NR^{13}R^{14}$, polyhalomethyloxy, $-NH-SO_2-R^8$, $-NH-SO_2-(C_{1-4}alkanediy)-CO-N(R^8)_2$; or Y is C_{1-6} alkyl substituted with cyano or with $-C(=O)R^8$;

(u) aryl is phenyl or phenyl substituted with one, two or three substituents each independently selected from halo, hydroxy, mercapto, C_{1-6} alkyl, hydroxy C_{1-6} alkyl, C_{1-6} alkyl $NR^{13}R^{14}$, C_{1-6} alkylcarbonyl, C_{1-6} alkyloxy, C_{1-6} alkyloxycarbonyl, C_{1-6} alkylthio, cyano, nitro, polyhalo C_{1-6} alkyl, polyhalo C_{1-6} alkyloxy, $-C(=O)-NR^{13}R^{14}$, R^7 or $-X_3-R^7$;

10 (v) Het is a monocyclic or bicyclic, partially saturated or aromatic heterocycle, wherein each of said carbocyclic or heterocyclic ring systems may optionally be substituted with one, two or three substituents each independently selected from halo, hydroxy, mercapto, C_{1-6} alkyl, hydroxy C_{1-6} alkyl, amino C_{1-6} alkyl, C_{1-6} alkylcarbonyl, C_{1-6} alkyloxy, C_{1-6} alkyloxycarbonyl, C_{1-6} alkylthio, cyano, nitro, polyhalo C_{1-6} alkyl, polyhalo C_{1-6} alkyloxy.

3. A compound according to claim 2 wherein all of the limitations (a) - (v) apply.

20

4. A compound according to claim 1 wherein one or more of the following limitations (a') - (v') apply :

(a') $-a^1=a^2-a^3=a^4-$ represents a bivalent radical of formula


25 $-CH=CH-CH=CH-$ (a-1);

(b') n is 1 or 2;

(c') m is 1 or 2;

(d') R^1 is hydrogen; C_{1-6} alkyl;

30 (e') each R^2 independently is hydroxy, halo, C_{1-6} alkyl optionally substituted with cyano or $-C(=O)R^6$, C_{2-6} alkenyl optionally substituted with cyano, C_{2-6} alkynyl optionally substituted with cyano, C_{1-6} alkyloxycarbonyl, carboxyl, cyano, nitro, amino, mono(C_{1-6} alkyl)amino, di(C_{1-6} alkyl)amino, $-S(=O)_pR^6$, $-NH-S(=O)_pR^6$, $-C(=O)R^6$, $-NHC(=O)H$, $-C(=O)NHNH_2$, $-NHC(=O)R^6$, $-C(=NH)R^6$ or a radical of formula



35 wherein each A_1 independently is N, CH or CR^6 ; and no more than two A_1 are N; A_2 is NH, O, S or NR^6 ;

-57-

- (f') X_1 is $-NR^5$ -, $-NH-NH$ -, $-N=N$ -, $-O$ -, $-C(=O)$ -, C_{1-4} alkanediyl, $-CHOH$ -, $-NR^{13}$ -, $C(=O)$ -, $-C(=O)-NR^{13}$ -, $-X_2-C_{1-4}$ alkanediyl- or $-C_{1-4}$ alkanediyl- X_2 -;
- (g') X_2 is $-NR^5$ -, $-O$ -;
- (h') R^3 is hydrogen, halo, C_{1-6} alkyl, $NR^{13}R^{14}$ -, $-C(=O)-NR^{13}R^{14}$ -, $-C(=O)-R^{15}$ -, $-X_3-R^7$;
- 5 C_{1-6} alkyl substituted with one or two substituents each independently selected from cyano, R^7 or $-C(=O)-NR^9R^{10}$; C_{2-6} alkenyl substituted with one or more substituents each independently selected from halo, cyano or $-C(=O)-NR^9R^{10}$; or C_{2-6} alkynyl substituted with one or more substituents each independently selected from halo, cyano, $-C(=O)-NR^9R^{10}$;
- 10 (i') X_3 is $-NR^5$ -or- O -;
- (j') R^4 is halo, hydroxy, C_{1-6} alkyl, C_{2-6} alkenyl, C_{2-6} alkynyl, C_{1-6} alkyloxy, cyano, nitro, $-C(=O)-NR^{13}R^{14}$ -, C_{1-6} alkyloxycarbonyl, C_{1-6} alkylcarbonyl, formyl, $-NR^{13}R^{14}$;
- (k') R^5 is hydrogen; C_{1-6} alkyl;
- (l') R^6 is C_{1-4} alkyl;
- 15 (m') R^7 is any of the specific monocyclic or bicyclic, partially saturated or aromatic carbocycles or monocyclic or bicyclic, partially saturated or aromatic heterocycles specifically mentioned in this specification, wherein each of said carbocyclic or heterocyclic ring systems may optionally be substituted with one, two or three substituents each independently selected from halo, hydroxy, mercapto, C_{1-6} alkyl,
- 20 hydroxy C_{1-6} alkyl, amino C_{1-6} alkyl, C_{1-6} alkylcarbonyl, C_{1-6} alkyloxy, C_{1-6} alkyloxycarbonyl, C_{1-6} alkylthio, cyano, nitro, polyhalo C_{1-6} alkyl, polyhalo C_{1-6} alkyloxy or aminocarbonyl;
- (n') R^8 is hydrogen or C_{1-4} alkyl;
- (o') R^9 and R^{10} each independently are hydrogen or C_{1-6} alkyl;
- 25 (p') R^{13} and R^{14} each independently are hydrogen or C_{1-6} alkyl;
- (q') R^{15} is C_{1-6} alkyl optionally substituted with cyano or $-C(=O)-NR^{13}R^{14}$;
- (r') R^{17} is cyano, halo, hydroxy, $-C(=O)-NR^{13}R^{14}$ -, C_{1-6} alkyl optionally substituted with cyano, $-C(=O)-NR^{13}R^{14}$ -, C_{2-6} alkenyl optionally substituted with cyano or $-C(=O)-NR^{13}R^{14}$ -, C_{2-6} alkynyl optionally substituted with cyano or $-C(=O)-NR^{13}R^{14}$ -, and,
- 30 where possible, R^{17} may also be attached to the $-b^1-b^2-b^3$ - moiety by a double bond whereby R^{17} is then $=O$, $=NH$, $=N-R^{15}$, $=N-R^7$, $=N-O-R^{15}$, $=N-O-R^7$, $=CH_2$, $=CH-C(=O)-NR^{13}R^{14}$ -, $=CH-R^7$, or $=CH-R^{15}$; wherein $=CH_2$ may optionally be substituted with cyano;
- (s') Q represents hydrogen or C_{1-6} alkyl or $-NR^9R^{10}$;
- 35 (t') Y represents hydrogen, hydroxy, halo, C_{1-6} alkyl, C_{1-6} alkyloxy, cyano, nitro, $NR^{13}R^{14}$ -, polyhalomethoxy, $-NH-SO_2-R^8$ -, $-NH-SO_2-(C_{1-4}$ alkanediyl)- $CO-N(R^8)_2$;
- (u') aryl is phenyl or phenyl substituted with one, two or three substituents each independently selected from halo, hydroxy, C_{1-6} alkyl, hydroxy C_{1-6} alkyl, C_{1-}

-58-

alkylcarbonyl, C₁₋₆alkyloxy, C₁₋₆alkyloxycarbonyl, C₁₋₆alkylthio, cyano, nitro, -C(=O)-NR¹³R¹⁴;

- (v') Het is a monocyclic or bicyclic, partially saturated or aromatic heterocycle, specifically mentioned in this specification, wherein each of said heterocyclic ring systems may optionally be substituted with one, two or three substituents each independently selected from halo, hydroxy, mercapto, C₁₋₆alkyl, hydroxyC₁₋₆alkyl, aminoC₁₋₆alkyl, C₁₋₆alkylcarbonyl, C₁₋₆alkyloxy, C₁₋₆alkyloxycarbonyl, C₁₋₆alkylthio, cyano, nitro, polyhaloC₁₋₆alkyl, polyhaloC₁₋₆alkyloxy.
- 5
- 10 5. A compound according to claim 4 wherein all of the limitations (a') - (v') apply.
6. A compound according to claim 1 wherein one or more of the following limitations (a'') - (v'') apply:
- 15 (a'') -a¹=a²-a³=a⁴- represents a bivalent radical of formula


$$-\text{CH}=\text{CH}-\text{CH}=\text{CH}- \quad (\text{a-1});$$
- (b'') n is 1;
- (c'') m is 1;
- (d'') R¹ is hydrogen; methyl;
- 20 (e'') R² is halo, C₁₋₆alkyl optionally substituted with cyano, C₂₋₆alkenyl optionally substituted with cyano, C₂₋₆alkynyl optionally substituted with cyano, C₁₋₆alkyloxycarbonyl, carboxyl, cyano, amino, mono(C₁₋₆alkyl)amino, di(C₁₋₆alkyl)amino;
- (f'') X₁ is -NR⁵-, -O-, -NR¹³-C(=O)-, -C(=O)-NR¹³-;
- 25 (h'') R³ is hydrogen, halo, C₁₋₆alkyl, NR¹³R¹⁴, -C(=O)-NR¹³R¹⁴, -C(=O)-R¹⁵; C₁₋₆alkyl substituted with cyano; C₂₋₆alkenyl substituted with cyano; or C₂₋₆alkynyl substituted with cyano;
- (j'') R⁴ is halo, hydroxy, C₁₋₆alkyl, C₂₋₆alkenyl, C₂₋₆alkynyl, C₁₋₆alkyloxy, cyano, nitro, -C(=O)-NR¹³R¹⁴, -NR¹³R¹⁴;
- 30 (k'') R⁵ is hydrogen; C₁₋₆alkyl;
- (m'') R⁷ is any of the specific monocyclic or bicyclic, partially saturated or aromatic carbocycles or monocyclic or bicyclic, partially saturated or aromatic heterocycles specifically mentioned in this specification, wherein each of said carbocyclic or heterocyclic ring systems may optionally be substituted with one, two or three substituents each independently selected from halo, hydroxy, mercapto, C₁₋₆alkyl, hydroxyC₁₋₆alkyl, aminoC₁₋₆alkyl, C₁₋₆alkylcarbonyl, C₁₋₆alkyloxy, C₁₋₆alkyloxycarbonyl, C₁₋₆alkylthio, cyano, nitro, polyhaloC₁₋₆alkyl, polyhaloC₁₋₆alkyloxy or aminocarbonyl;
- 35

- (n") R⁸ is hydrogen or C₁₋₄alkyl;
 (o") R⁹ and R¹⁰ are hydrogen;
 (p") R¹³ and R¹⁴ are hydrogen;
 (q") R¹⁵ is C₁₋₆alkyl optionally substituted with cyano;
 5 (r") R¹⁷ is cyano, -C(=O)-NR¹³R¹⁴, C₁₋₆alkyl optionally substituted with cyano, -C(=O)-NR¹³R¹⁴; C₂₋₆alkenyl optionally substituted with cyano or -C(=O)-NR¹³R¹⁴; C₂₋₆alkynyl optionally substituted with cyano or -C(=O)-NR¹³R¹⁴; and, where possible, R¹⁷ may also be attached to the -b¹-b²-b³- moiety by a double bond whereby R¹⁷ is then =O, =NH, =N-R¹⁵, =N-R⁷, =N-O-R¹⁵, =N-O-R⁷, =CH₂, =CH-C(=O)-
 10 NR¹³R¹⁴, =CH-R⁷, or =CH-R¹⁵; wherein =CH₂ may optionally be substituted with cyano;
 (s") Q represents hydrogen or -NR⁹R¹⁰;
 (t") Y represents hydrogen, hydroxy, halo, C₁₋₆alkyl, C₁₋₆alkyloxy, cyano, NR¹³R¹⁴, -NH-SO₂-R⁸, -NH-SO₂-(C₁₋₄alkanediyl)-CO-N(R⁸)₂;
 15 (u") aryl is phenyl or phenyl substituted with one, two or three substituents each independently selected from halo, hydroxy, C₁₋₆alkyl, C₁₋₆alkyloxy, C₁₋₆alkylthio, cyano, nitro;
 (v") Het is a monocyclic or bicyclic, partially saturated or aromatic heterocycle, specifically mentioned in this specification, wherein each of said heterocyclic ring
 20 systems may optionally be substituted with one, two or three substituents each independently selected from halo, hydroxy, mercapto, C₁₋₆alkyl, hydroxyC₁₋₆alkyl, aminoC₁₋₆alkyl, C₁₋₆alkylcarbonyl, C₁₋₆alkyloxy, C₁₋₆alkyloxycarbonyl, C₁₋₆alkylthio, cyano, nitro, polyhaloC₁₋₆alkyl, polyhaloC₁₋₆alkyloxy.
- 25 7. A compound according to claim 6 wherein all limitations (a") - (v") apply.
8. A compound of formula (I) as claimed in claims 1 - 7 for use as a medicine.
9. Use of a compound of formula (I) as claimed in claims 1 - 7 in the manufacture of a
 30 medicament for the treatment or prophylaxis of an infectious disease.
10. A pharmaceutical composition comprising (a) an effective amount of a compound of formula (I) as claimed in claim 1, and (b) pharmaceutically acceptable carrier.
- 35 11. A process for preparing a composition as claimed in claim 10 comprising mixing the compound of formula (I) with the carrier.


-60-

12. A product comprising (a) a compound of formula (I) as claimed in claims 1 - 7, and (b) another antiretroviral compound as a combined preparation for simultaneous, separate or sequential use in the treatment or prophylaxis of HIV infection.
- 5 13. A process for preparing a compound of formula (I) as claimed in claims 1 - 7, characterized in that


(a) an indane of formula (a-1) is reacted with a reagent (a-2):



reagent (a-2) being of general formula



- 10 The substituents have the meanings specified in claims 1-8, X_1 and W_1 are selected such that a linking radical X_1 is performed;
- (b) reacting a starting material (a-10) by reaction with an amino substituted aromatic compound (a-11) in an arylation type of reaction:



and wherein the aryldane (a-10) is obtainable by reacting an indane of formula (a-1) with a pyrimidine (a-9) wherein the substituents have the meanings specified in claims 1-

5 8, and X_1 and W_1 are selected such that a linking radical X_1 is formed;

(c) and if desired converting the compounds of formula (I) into each other by a suitable conversion reaction;

10 (d) and if further desired preparing a a pharmaceutically acceptable addition salt, a quaternary amine thereof; or a stereochemically isomeric form thereof.

INTERNATIONAL SEARCH REPORT

International Application No
P2004/050175

A. CLASSIFICATION OF SUBJECT MATTER

IPC 7 C07D239/47 A61K31/505 C07D239/48 C07D413/12 A61P31/18

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

IPC 7 C07D A61K A61P

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the International search (name of data base and, where practical, search terms used)

EPO-Internal, CHEM ABS Data

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	WO 99/50256 A (KOYMANS LUCIEN MARIA HENRICUS ; HEERES JAN (BE); DAEYAERT FREDERIK) 7 October 1999 (1999-10-07) example 84	1-13
X	WO 02/36578 A (BONHAM LYNN ; KLEIN J PETER (US); LEUNG DAVID W (US); TANG NORINA M) 10 May 2002 (2002-05-10) example 10	1
Y	EP 0 834 507 A (JANSSEN PHARMACEUTICA NV) 8 April 1998 (1998-04-08) table 5	1-13
	----- -/-	

☒ Further documents are listed in the continuation of box C.

☒ Patent family members are listed in annex.

* Special categories of cited documents:

- *A* document defining the general state of the art which is not considered to be of particular relevance
- *E* earlier document but published on or after the international filing date
- *L* document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)
- *O* document referring to an oral disclosure, use, exhibition or other means
- *P* document published prior to the international filing date but later than the priority date claimed

- *T* later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention
- *X* document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone
- *Y* document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.
- *&* document member of the same patent family

Date of the actual completion of the international search

15 June 2004

Date of mailing of the international search report

24/06/2004

Name and mailing address of the ISA

European Patent Office, P.B. 5818 Patentlaan 2
NL - 2280 HV Rijswijk
Tel. (+31-70) 340-2040, Tx. 31 651 epo nl,
Fax (+31-70) 340-3016

Authorized officer

Lauro, P

INTERNATIONAL SEARCH REPORT

International Application No
PCT/EP2004/050175

C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
Y	WO 99/50250 A (AKEN KOEN JEANNE ALFONS VAN ;KOYMANS LUCIEN MARIA HENRICUS (BE); H) 7 October 1999 (1999-10-07) cited in the application the whole document	1-13
Y	WO 00/27825 A (AKEN KOEN JEANNE ALFONS VAN ;KOYMANS LUCIEN MARIA HENRICUS (BE); H) 18 May 2000 (2000-05-18) cited in the application the whole document	1-13

INTERNATIONAL SEARCH REPORT

 Int. Application No
 PCT/EP2004/050175

Patent document cited in search report		Publication date	Patent family member(s)	Publication date
WO 9950256	A	07-10-1999	AU 758624 B2	27-03-2003
			AU 3599799 A	18-10-1999
			BG 104716 A	30-04-2001
			BR 9909197 A	05-12-2000
			CA 2324921 A1	07-10-1999
			CN 1295566 T	16-05-2001
			EE 200000535 A	15-04-2002
			WO 9950256 A1	07-10-1999
			EP 1066269 A1	10-01-2001
			HR 20000621 A1	30-04-2001
			HU 0101372 A2	28-10-2001
			ID 26043 A	16-11-2000
			JP 2002511390 T	16-04-2002
			NO 20004809 A	24-11-2000
			NZ 506787 A	28-02-2003
			PL 343195 A1	30-07-2001
			SK 14052000 A3	11-06-2001
			TR 200002761 T2	22-01-2001
			US 2002115668 A1	22-08-2002
			US 6150360 A	21-11-2000
			US 6372729 B1	16-04-2002
			EP 0945447 A1	29-09-1999
			ZA 200006042 A	26-10-2001
WO 0236578	A	10-05-2002	AU 1665002 A	15-05-2002
			WO 0236578 A2	10-05-2002
			US 2003100557 A1	29-05-2003
			US 2002103195 A1	01-08-2002
EP 0834507	A	08-04-1998	AP 914 A	18-12-2000
			AU 740809 B2	15-11-2001
			AU 3926697 A	09-04-1998
			BR 9704937 A	06-06-2000
			CA 2216486 A1	01-04-1998
			CN 1180698 A ,B	06-05-1998
			CZ 9702993 A3	11-11-1998
			EE 9700253 A	15-04-1998
			EP 0834507 A1	08-04-1998
			HR 970526 A1	31-10-1998
			HU 9701596 A2	28-06-1999
			ID 19599 A	23-07-1998
			IL 121849 A	26-08-2001
			JP 10114759 A	06-05-1998
			NO 974368 A	02-04-1998
			NZ 328854 A	27-10-2000
			OA 10620 A	03-09-2002
			PL 322369 A1	14-04-1998
			RU 2186774 C2	10-08-2002
			SG 53075 A1	28-09-1998
			SK 131997 A3	11-06-1999
			TR 9701070 A2	21-04-1998
			TW 411335 B	11-11-2000
			US 2002147181 A1	10-10-2002
			US 2003199473 A1	23-10-2003
			US 6380194 B1	30-04-2002
			ZA 9708766 A	30-03-1999
WO 9950250	A	07-10-1999	AT 232521 T	15-02-2003

INTERNATIONAL SEARCH REPORT

International Application No
PL/EP2004/050175

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
WO 9950250 A		AU 751573 B2	22-08-2002
		AU 3599699 A	18-10-1999
		BG 104738 A	30-04-2001
		BR 9909191 A	05-12-2000
		CA 2324919 A1	07-10-1999
		CN 1295564 T	16-05-2001
		DE 69905306 D1	20-03-2003
		DE 69905306 T2	27-11-2003
		DK 945443 T3	02-06-2003
		EA 2973 B1	26-12-2002
		EE 200000532 A	15-02-2002
		WO 9950250 A1	07-10-1999
		EP 1245567 A1	02-10-2002
		EP 0945443 A1	29-09-1999
		ES 2193660 T3	01-11-2003
		HR 20000620 A1	30-06-2001
		HU 0101204 A2	28-10-2001
		ID 26291 A	14-12-2000
		JP 3507917 B2	15-03-2004
		JP 2002509920 T	02-04-2002
		NO 20004810 A	26-09-2000
		NZ 506679 A	26-11-2002
		PL 343196 A1	30-07-2001
		PT 945443 T	30-06-2003
		SI 945443 T1	31-08-2003
		SK 14062000 A3	11-06-2001
		TR 200002760 T2	21-12-2000
		TW 531534 B	11-05-2003
		US 2003083317 A1	01-05-2003
		US 6197779 B1	06-03-2001
		US 2001011094 A1	02-08-2001
		EP 0945442 A1	29-09-1999
		ZA 200006044 A	26-10-2001
WO 0027825 A	18-05-2000	AT 233740 T	15-03-2003
		AU 762523 B2	26-06-2003
		AU 6200899 A	29-05-2000
		BG 105418 A	30-11-2001
		BR 9915552 A	14-08-2001
		CA 2350801 A1	18-05-2000
		CN 1322198 T	14-11-2001
		CZ 20011533 A3	17-10-2001
		DE 69905683 D1	10-04-2003
		DE 69905683 T2	18-03-2004
		DK 1002795 T3	30-06-2003
		EA 4049 B1	25-12-2003
		EE 200100252 A	15-10-2002
		WO 0027825 A1	18-05-2000
		EP 1270560 A1	02-01-2003
		EP 1002795 A1	24-05-2000
		ES 2193664 T3	01-11-2003
		HK 1025330 A1	13-06-2003
		HR 20010161 A1	28-02-2002
		HU 0104177 A2	28-03-2002
		ID 28376 A	17-05-2001
		JP 2002529456 T	10-09-2002
		NO 20011696 A	04-04-2001
		NZ 511116 A	29-08-2003

INTERNATIONAL SEARCH REPORT

Int. Patent Application No
PCT/EP2004/050175

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
WO 0027825	A	PL 347586 A1	08-04-2002
		PT 1002795 T	31-07-2003
		SI 1002795 T1	31-10-2003
		SK 6032001 A3	07-01-2002
		TR 200101306 T2	22-10-2001
		US 2003114472 A1	19-06-2003
		US 2004039005 A1	26-02-2004
		ZA 200103769 A	12-08-2002